

Title :	<b>AN 146</b>	<b>Bleeding Relief and repair</b>	Pages: 1
Doc Name :	<b>For all species</b>		Related Products :
Last Updated:	12.01.17		

#### HOMEOPATHIC MEDICINE

**One dose** = approx. 2 to 3 pills orally in gum of mouth or crush pills (no need to open mouth) or 3 to 5 drops on body/skin as one dose. If formula is in liquid do not put this in mouth as it contains alcohol.  
Refer to: **Bottle label** for suggested frequency repeat dosing.

#### Extra help:

To help strengthen blood vessels as well as assisting iron absorption (if anemic) use **Vitamin C**

**You will need:** One (1) Oral syringe (without the needle) (from chemist)

Vitamin C tablets (crush the Vit C tablets if you have this at home)

or

(if you need to buy vit c then best to buy the "crystal" powder as it dissolves well in fluids)

Example: BUY A packet of **Sodium Ascorbate** Vitamin C crystal (white powder)

#### THERAPY (A) **Serious/ life threatening**

- **CAT / small dog / any small animal** - mix 3 teaspoons of crystals or powdered vitamin C in 1 cup of warm filtered water and give approx. 5mls – 10mls orally - every 15 minutes for up to 4 hours, ( approx 20 oral doses) then reduce to every 3 - 4 hours on improvement. Then stop when back to normal self.
- **DOGS /horse /cow / sheep etc** - mix 5 teaspoons of powder in 1 cup of warm water and give 20ml every 15 minutes up to 4 hours, reduce frequency on improvement to every couple of hours, then stop when back to normal.

#### THERAPY (B) **Semi Serious**

- **CAT / small dog / any small animal** - mix 3 teaspoons of crystals or powdered vitamin C in 1 cup of warm filtered honey water or goats milk or chicken broth.  
\*Using the oral syringe: draw up liquid mix and give approx. 5mls orally (side of mouth)- every 1 to 2 hours – 4 times a day. On improvement, can reduce frequencies per day. Then can stop when back to better health.
- **DOG / horse. Sheep/ cow** - mix 5 teaspoons of powder in 1 cup of warm filtered honey water or goats milk or chicken broth.

\*Using Oral syringe draw up mixture and give approx. 20ml orally (side of mouth) every 1 hour for 4 hours hours. Then repeat again following days. If no improvement (or loose stools seen) within 2 days,

Then up the amount given for each dose. i.e. 40ml every 2 hours 4 times daily for another 2 to 3 days. When improvement seen (or loose stools) you can then slightly reduce the amount each time and then maintain.