

|   |                  |
|---|------------------|
| <p><b>Product Code AN110 &amp; 42 &amp; 338</b></p> <p><b>Chemical Poisoning Detox Set</b></p> <p>Chemical Drugs Side effects and Damages toxicities poisonings or<br/>fatal allergy reactions</p> <p><b>Last Updated: 24-2-21</b> <span style="float: right;"><b>For all species &amp; ages</b></span></p> | <p>71 Pages:</p> |
|---|------------------|

\* THIS IMPORTANT INFORMATION IS ABOUT WHAT CHEMICAL DRUGS CAN DO TO THE BODY AND SYMPTOMS TO LOOK OUT FOR. IF YOU ARE CURRENTLY GIVING YOUR DOG OR CAT ANY OF THE BELOW CHEMICAL PRODUCTS. PLEASE CONSIDER DETOX YOUR PET AND WHERE POSSIBLE USE SAFE ALTERNATIVES. **The found of 'Homeopathic medicine' physician Samuel Hahnemann said to "remove the obstacles to the cure" so detox of these chemical is vital.**

## HOMEOPATHIC DETOX

HAMPL Chemical Poisoning 110 30ml or HAMPL Snail Pellet 338 30ml plus  
Activated Charcoal powder , Vitamin C powder  
Or HAMPL Pet Cleanse 42

\* This one is good for not well since taking a drug, or insecticide products, now seizures, paralysed, High liver enzymes etc. Or we can provide the one drug antidote instead of the combination detox.

~ Also use the formulas for the symptoms of damages... seizure - neurological damages repair, paralysis etc ask out practitioners our recommendation or just a detox using [HAMPL Pet Cleanse 042 30ml plus Activated Charcoal powder \(set of 2\)](#)

This one is good for pets that have not had a lot of drugs or other dangerous chemicals but still need cleansing. High liver enzymes etc.

\* (asterisk) means we have this in homeopathic form of the toxic drug is in the above formulas to aid a detox from this drug.

On request we can provide just one drug detox specifically to the drug your pet has not been well since taking it or had acute side effects from within minutes or hours. If you dont see a drug that your pet is taking or did take, please send a tiny sample of it to our clinic and we will post out the detox homeopathic version to you

## WHAT TO DO STRAIGHT AWAY

[ANTI DOTING poisons](#) using Vitamin C *\*See page 65 Instructions*  
With Activated Charcoal – *to absorb poisons in gut.* *\* See page 73*

**Note** ALL chemical products & drugs affect the liver and some also affect (damage) kidneys as well.

## Page Content Index

|   |                |
|---|----------------|
| COMFORTIS® (spinosad) Or Simparica (sarolaner Chewable)                   | - Page 6       |
| MILBEMAX® Drug  | - Page 7       |
| PANACUR® Liquid Or Purina TotalCare or Drontal or Virbantel               | - Page 7       |
| ADVOCATE® Drug (Advantage multi)  | - Page 8       |
| NEXGARD® Drug   | - Page 16 - 17 |
| HEARTGUARD Plus, HeartGuard, Nuheart - IVERMECTIN® P                      | - Page 18 - 19 |
| FRONTLINE® Fipronil (frontline gold, frontline plus)                      | - Page 20 - 21 |
| TRIFEXIS® (spinosad + milbemycin oxime)                                   | - Page 22      |
| SIMPATICO TRIO  | - Page 23      |
| BRAVECTO tablets  | - Page 24 - 27 |
| SENTINEL® tablets & The Big 5 Pack <b>Milbemycin oxime (Interceptor®)</b> | - Page 28 - 29 |
| PROFENDER®  | - Page 30      |
| REVEOLUTION Plus®   | - Page 30      |
| AMITRAZ®  | - Page 31      |
| SERESTO Collars   | - Page 32      |
| APOQUEL® Drug   | - Page 32 - 34 |
| MALASEB®  | - Page 35      |
| ANTIROBE®   | - Page 36 - 37 |
| VERAFLOX® Pradofloxacin   | - Page 37      |
| METRONIDAZOLE®  | - Page 37      |
| DOXYCYCLINE®  | - Page 38      |
| CLAVAMOX® Amoxy Clavulanic acid   | - Page 38      |
| CONVENIA®   | - Page 39      |
| BAYTRIL® (Enrofloxacin)   | - Page 39      |
| FLUCONAZOLE® Drug   | - Page 40      |
| ELOXORAL® Drug Metecam or Meloxicam                                       | - Page 45      |
| TRAMADOL® Drug  | - Page 46      |
| <b>BUPRENORPHINE® (Buprenex)</b>  | - Page 46      |
| FIROXIB® Drug   | - Page 47      |
| SOMEPRAZOLE® Drug   | - Page 48 - 49 |
| PREDNISON® Drug ( <b>Prednisolone</b> )                                   | - Page 49 - 50 |
| PIMOBENDAN® Drug ( <b>Vetmedin</b> )                                      | - Page 51      |
| ENALPRIL MALEATE® Drug  | - Page 52      |
| PROIN® Drug ( <b>Phenylpropanolamine</b> )                                | - Page 53      |
| PROPALIN® Drug or <b>Urex Drug (UTI)</b>                                  | - Page 54      |
| UREX Drug (urinary) or Piroxicam (bladder cancer drug)                    | - Page 55      |
| GABAPENTIN® Drug  | - Page 56 - 59 |
| Iron EDTA, Metaldehyde and Methiocarb e.g <b>Snail &amp; Slug Baits</b>   | - Page 60      |

*The Big 5 Pack \* Interceptor Spectrum: Milbemycin oxime,  
Praziquantel \* Credelio: Lotilaner*

# INSECTICIDE POISON

\* **RoundUp (glyphosate) "used as a weed killer"** (and human-pet killer) The **carcinogenic** insecticide spray which is commonly used around parks, gardens at home worldwide. This highly toxic chemical is made by Monsanto. **The Lawsuit** filed against them for over 4,000 across USA diagnosed with cancer.

<https://www.baumhedlundlaw.com/toxic-tort-law/monsanto-roundup-lawsuit/>

## TEST kit

**Australia or Canada testing yourself or pets**

<http://hrilabs.org/animalstudy/>

**USA** <https://shop.mercola.com/product/2057/1/glyphosate-environmental-exposure-test-kit>

**Glyphosate homeopathic 30C is in the Pet Cleanse 42 and Chemical Detox 110 homeopathic formula.**

Optional to test for it, but as its been used in gardens and parks we could be guaranteed we have been affected by it. You and your pets (farm animals) can as safely use the homeopathic detox combination pet cleanse or chemical detox. Or alternatively can request it on its on.

All gentle detoxing in conjunction with activate charcoal powder. Recently we sent this detox for a cat with cancer, within a 3 days her cat was more energetic and eating eagerly and started putting on weight again.

This lady lived on a farm and knew farmers using the spray miles away. Its a carcinogenic!

## SOME OF THE VETERINARY DRUGS OR OVER THE COUNTER FROM PET SHOPS THAT are given to our pets.

\* Side effects after dogs eating...Nestle Purina's multicolored kibble

<https://www.dogsnaturallymagazine.com/fda-releases-report-toxic-ingredients-beneful/>

### **THE BELOW PRODUCTS, SPRAYS, DROPS, TABLETS, INJECTIONS FROM 1 TO 5**

are classified as a insecticide poison. Which is a Neurotoxin to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.

# INSECTICIDE POISON - Pet Products

FDA Alerts Pet Owners and Veterinarians About Potential for Neurologic Adverse Events with chemical Flea and Tick Products. <https://wjla.com/news/nation-world/fda-warns>

## Physical Symptoms of Neurological Problems

Physical symptoms of neurological problems may include the following:

- Partial or complete paralysis
- Muscle weakness
- Partial or complete loss of sensation
- Seizures
- Poor cognitive abilities
- Unexplained pain
- Decreased alertness
- Fatal

## **ALERT** Fact Sheet for Pet Owners and Veterinarians about Potential Adverse Events Associated with

### **Isoxazoline Flea and Tick Products**

Content current as of: 07/31/2020 Fast Facts

The FDA is alerting pet owners and veterinarians of the potential for neurologic adverse events in dogs and cats when treated with drugs that are in the isoxazoline class. The drugs in this class are Bravecto (fluralaner) tablets for dogs, Bravecto (fluralaner) topical solution for cats and dogs, Bravecto Plus (fluralaner and moxidectin) topical solution for cats, Bravecto 1-month (fluralaner) tablets for cats, Nexgard (afoxalaner) tablets for dogs, Simparica (sarolaner) tablets for dogs, Simparica Trio (sarolaner, moxidectin and pyrantel) tablets for dogs, Credelio (lotilaner) tablets for dogs and cats, and Revolution Plus (selamectin and sarolaner) topical solution for cats. These products are approved for the treatment and prevention of flea infestations, and the treatment and control of tick infestations. Some of these products also have other indications. Although these products can and have been safely used in the majority of dogs and cats, pet owners should consult with their veterinarian to review their patients' medical histories and determine whether a product in the isoxazoline class is appropriate for their pet.

## *What should I know?*

*The FDA considers products in the isoxazoline class to be safe and effective for dogs and cats but is providing this information so that pet owners and veterinarians can take it into consideration when choosing flea and tick products for their pets.*

~ **Isoxazoline** products have been associated with **neurologic adverse** reactions, **including muscle tremors, ataxia, and seizures in some dogs and cats;**

~ **seizures may occur** in animals without a prior history;

### ***What products are in the isoxazoline class?***

- The FDA- drugs in this class are
  - Bravecto (fluralaner) tablets for dogs
  - Bravecto (fluralaner) topical solution for cats and dogs
  - Bravecto Plus (fluralaner and moxidectin) topical solution for cats
  - Bravecto 1-month (fluralaner) tablets for cats
  - Credelio (lotilaner) tablets for dogs and cats
  - Nexgard (afoxaloner) tablets for dogs
  - Simparica (sarolaner) tablets for dogs
  - Simparica Trio (sarolaner, moxidectin and pyrantel) tablets for dogs
  - Revolution Plus (selamectin and sarolaner) topical solution for cats

<https://www.fda.gov/animal-veterinary/animal-health-literacy/fact-sheet-pet-owners-and-veterinarians-about-potential-adverse-events-associated-isoxazoline-flea>

# FLEA Insecticide products

## \* COMFORTIS Comfortis® (spinosad) flea treatment.

Is classified as a **insecticide poison**. A **Neurotoxin** to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.

Spinosad contains two major factors, spinosyn A and spinosyn D.

**FOR CATS:** Adverse reactions were: *lethargy, decreased appetite, weight loss, and diarrhea. Use with caution with concomitant extra-label use of ivermectin. Avoid Comfortis in breeding, pregnant, or lactating cats.*

**FOR DOGS:** Adverse reactions reported in decreasing order of frequency are: *vomiting, depression / lethargy, decreased appetite, incoordination, diarrhea, itching, trembling, excessive salivation, trembling / twitching, salivation / drooling, seizures, ataxia, mydriasis, blindness and disorientation, excessive dilation of pupils, blindness and disorientation.*

**With even more serious adverse reactions have been reported following use of ivermectin with Comfortis.:** *acute severe vomiting, lethargy, anorexia, ataxia, diarrhea, pruritus, trembling in pain and nerve damage, hypersalivation and seizures, (death within a day of use).* CHEWABLE TABLETS had a much higher percentage of acute adverse side effects compared to Active Topical drug.

## Simparica (sarolaner Chewable) Flea and Tick

Simparica works by impacting the nervous system and is a **classified as a insecticide poison**.

A **Neurotoxin** to be precise The most frequently reported adverse reactions were vomiting and diarrhea. Simparica may cause abnormal neurologic signs such as **tremors, decreased conscious proprioception (awareness of body position and movement), ataxia (loss of full control of bodily movements), decreased or absent menace blinking reflex or seizures.**

### HEALTH NEWS Flea, Tick pills causes nerve (seizures) reaction in pets, FDA warns

**Sept. 21, 2018, 4:34 AM AWST By Maggie Fox**

Pills or chews that protect dogs and cats against fleas can cause neurological problems such as stumbling or seizures, and pet owners need more warning, the Food and Drug Administration said Thursday. The FDA **issued an alert** to owners and veterinarians and said it was requiring clearer labels on the products. *The flea pills and chews all contain a pesticide called isoxazoline, the FDA said. They include products sold under the brand names Bravecto, Nexgard and Simparica. "Another product in this class, Credelio, recently received FDA approval. These products are approved for the treatment and prevention of flea infestations, and the treatment and control of tick infestations," the FDA said in a statement.*

<https://www.nbcnews.com/health/health-news/flea-tick-pills-can-cause-nerve-reactions-pets-fda-warns-n911536>

# WORMERS

## \* **PANACUR liquid Wormer**

**A worming drug - which is classified as an insecticide poison.**

A Neurotoxin to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function. Panacur Liquid for Cats and dog , horse and other animals. Fenbendazole, marketed under the brand name Panacur, is a common anthelmintic -or dewormer - for cats, dogs, horses and other animals. SIDE EFFECTS - The FREQUENT side effects that may occur include infrequent vomiting or diarrhea.

## **Purina TotalCare or Drontal or Virbantel "Wormer"**

**A worming drug - which is classified as an insecticide poison.**

Chemical insecticide poison of - Pyrantel embonate and Niclosamide, pyrantel pamoate/praziquante  
**Pyrantel embonate side effects:** Skin rash, Abdominal or stomach cramps and pain causing nausea, vomiting, diarrhea, dizziness, drowsiness, headache Irritability, loss of appetite, nausea or vomiting, trouble in sleeping Cat Side effects of Drontal - loss of appetite, drooling (in cats), vomiting, lethargy (lack of energy), and diarrhea. Dog and puppy Side effects of Drontal - suffer from such as lack of appetite, gastrointestinal upset. **The** lining of the intestines become inflamed, causing diarrhea, **vomiting** and lack of appetite  
**Niclosamide side effects all pets:** Abdominal or stomach cramps or pain, diarrhea, loss of appetite, nausea or vomiting, dizziness or lightheadedness, drowsiness confusion, skin rash

## \* **MILBEMAX**

(wormer and heartworm drug) which is classified as an insecticide poison. A Neurotoxin to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.

**Milbemat** is a toxic insecticide drug used to control of gastrointestinal worms in pets including roundworm (*Toxocara cati*, *Toxascaris leonina*), hookworm (*Ancylostoma tubaeforme*) and tapeworm (*Dipylidium caninum*, *Taenia spp.*) and heartworm What are the active ingredients in Milbemax DRUG and how do they work? Milbemax contains praziquantel and milbemycin. **Both ingredients act by disrupting the nervous system of the worms resulting in their paralysis and death.** **SIDE EFFECTS - Systemic signs (such as lethargy), neurological signs (such as ataxia and muscle tremors) and / or gastrointestinal signs (such as emesis and diarrhoea) have been observed after administration of the veterinary medicinal product.**

COMMENTS from consumers

Hi - I have a 14 year old cat that has had a very bad reaction to Milbemax wormer he had it administered on Tuesday morning and within hours: developed neurological symptoms, falling over. he has gone deaf ( he was not before) and will not drink or eat. Now its Friday and he still has no appetite and will not touch any food though he has been going in garden and may be drinking from bird bath. have been back to vet and she cant offer any suggestions, they have spoken to Novartis who say have not had any problems but reading this forum and other internet news these side effects are not uncommon. He is now steady on his feet but still will not eat, vet has checked throat and swallowing seems to be fine. I have managed to put a few flakes of white fish in his mouth and he will swallow. he seems interested in food but picks it up and drops it. (normally he would eat three sachets of food a day)

# Skin conditions

## \* ADVOCATE Drug (Advantage multi)

which is classified as an insecticide poison. A **Neurotoxin to be precise.** These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.

### What is Advocate used for?

Advocate is used for cats, ferrets and dogs suffering from, or at risk from, mixed parasitic infections (caused by several different types of parasites): For the treatment and prevention of flea, flea allergy dermatitis, ear mite, lice, sarcoptic mange and demodicosis, heartworm. The two active ingredients are: ~ **Imidacloprid & Moxidectin.** The ingredient, **Imidacloprid acts as a neurotoxin,** causing parasites to lose control of their central nervous system. E.g *spasm and seizures.*

### Side Effects and Precautions - Advantage - Advocate products

Drooling, depression, weakness, muscle spasms, dilated pupils, and panting.

If your pet shows any adverse symptoms, then you should speak to your veterinarian straight away and wash the treated area with warm water and a pet-safe shampoo.

### Here is a list of the probable side effects and what to watch for in your pet:

- Facial swelling
- Hives
- Scratching
- Diarrhea
- Vomiting
- Shock
- Seizures
- Pale gums
- Cold limbs
- Coma and death

### COMMENTS from consumers

Reference: <https://thedoginfoblog.com/advocate-for-dogs>

#### Linda

my friends dog was given this for fleas. he is now suffering and the vets have no idea whats wrong with him. They seem to think its a neurological problem. Dog cant use its back legs cant stand up and is currently still in the vets on a drip and being sedated. They have done tests but so far nothing showed up and waiting for more tests. He's been in now for 4 days. They only thing my friend did was give him some of this. reason puppy was taken to vets ( hes 8 months old) is he couldnt get up, whne he did he fell back down as if legs gave way. Vet ( PDSA) him some injections and dog got worse on the way home and was taken back on saturday, we are now wednesday and no closer to knowing exactly what is wrong. My friend is so upset and worried that if it is neurological that the puppy may not get better. he is thinking the worst at the moment. After searching the internet for him i came up with this site and notice that it does mention side effects. do you know if there would be any cure as its killing us not knowing what is wrong with him or what may have caused it.

But taking this medication is the only thing the owner can think of. Dog is at the PDSA as my friend does not work and is on benefits, we are just hoping the dog is in good hands. Dog is a small terrier x

**Joanna M.**

Hi. My dog has been having seizures. I am convinced that it is Advantage Multi as well. I would also take legal action. **Angela**

I also don't know how old this post is, but at least I can give you some feedback that may help. I work in a homeopathic clinic, for humans not veterinary and one of our patients came in and said their dog had collapsed after going outside after they had sprayed insecticide – I know it isn't exactly the same but it doesn't matter from homeopathic point of view. The vet had been saying that probably the dog wouldn't survive and they were very upset.

Our Homeopath gave them something for their dog, based on the symptoms. Within half an hour he was up and running around and totally different. Of course the family were overjoyed. If you can find a decent homeopath where you are they can probably help you. I live in the Middle East so not able to suggest homeopath for other countries, but you may also find online.



**STELLA MARIE**

My baby tucker too, he suffering from neurological disorder right now. Can't walk straight and keep on crying. I noticed it after 4 days of advocate treatment. I'm very very sad



**Roxanne**

My 14 year old Pomeranian poodle mix received his first dose of advantagemulti on Oct 23 and is having neurological symptoms. He acts startled, disoriented, is lethargic at times, his tail is down some of the time and he will just jump of the bed, couch etc. as if something is attacking him. He is a very active, healthy 14 year old, and the onset of these symptoms have to be related to the drug.

My vet came to see him today and agrees with me that it was probably the advantage multi. I will not use it again, and Am praying that he gets better as the drug gets out of his system. Years ago a Bayor product caused a serious illness ifor my mother-in-law and the medicine was taken off the market. They seem a little too quick for profit to do the appropriate testing and disclosure. Let me now if you have gotten any response from Bayor. Thank you



**Samantha Lyons**

I'm taking my dog to the vets in an hour. He isn't walking properly, is very unbalanced and sounds the same as some of these dogs on here. I treated him with advocate a couple of hours ago my poor boy is really going through it of late.



**Linda**

Hi I've just read some of your remarks. I have an old British bulldog. I was told to treat him with advocate when I collected him. Which I have been. I know these dogs have symptoms. But he's only 9 months and acts like he's got arthritis. I have taken him to the vet they think I'm walking him too much.



**Kim**

My French bulldog has exactly same reaction pdsa vets kept selling more saying advocate ok. They so wrong my dog now very ill



**Amy Heighton**

Hi Katherine, I don't know how old this post is but my 7 month old dog is experiencing the same symptoms. Please get in touch with me as I am very concerned that it is Advantage that has caused this. Thanks, Amy



**Kim**

Defo advantage I've used it 4 times symptoms getting worse. Never again



**Brian Jay**

My dog a border collie cross. Lost control of back legs. I put her down. Do you have a lawyer? This cant happen to anyone else. My dog was so sweet.



**Judy**

My collie is a year old and been having this treatment every month, now he really hates it to the point of biting us if we try to hold him for it, and the dislike is really bad, I've never seen a flea on him and now wondering whether to continue with it.



**Chris** - It should never be used on collies. It says so on the instructions.



**J**

Hi, my dog too had same reaction Sunday morning on advocate,back legs kept going,swaying all over the place,never again.x



**Karen**

We have just had a similar experience after using Advocate. It was terrifying. . The strange thing is we been using for two years . [karenknight269@gmail.com](mailto:karenknight269@gmail.com)



**Evelyne phelan**

Been giving this to my dog chihuahua for 2 years n dog has developed liver bile problem am not sure if it's this but dog has been very sick bloods shower liver problem n I wonder if this drug is toxic for her liver.



**Chris B**

Sorry about your dog ,get the vet to check for SRMA (steroid responsive meningitis ) it is quite common. Dog hurts to move ,neck pain, stiff joints not willing to move the only way to diagnose properly if a spinal tap/ MIR scan. My dog suffered with this and had a relapse on two occasions 3 days after Advacate , Prednisalone tabs work like magic within 4 hours you would not know he'd been ill.

If he responds he'll have to be on the steroids a good while and slowly weaned off. Hope this helps.



**Katherine**

Hi Linda The exact same thing happened to my dog, Speckles. I am sad to tell you that you will not get any support from Bayer (financial or otherwise). They will insist to the end that it was not their product. Vets will be equally hesitant to condemn the product. I had all the tests run on Speckles – MRI, X-ray, Genetics, etc. and no vet has ever discovered an underlying problem. Your puppy can live, but it is going to be a very hard road, and it might take a very long time. This product has a 35 day half life. This means it is going to take 35 days for the product to reduce by 50% in the dog's system. Speckles could not walk for about 2 months. You will want to begin to control for secondary issues such as urinary tract infections and megacolon. Ask if you can go ahead and use an antibiotic as a urinary tract protective. Find the best internal medicine vet specialist in your area and go there ASAP. Emergency vets are also pretty open to these types of reactions because they have seen them before. Please email me at [savespeckles@gmail.com](mailto:savespeckles@gmail.com) or [katayoe@gmail.com](mailto:katayoe@gmail.com). I hope to find enough people who this has happened to so we can form a class action lawsuit. Best wishes to you and your dog. Speckles will be sending good vibes!

Here is Speckles' story: <https://www.youtube.com/watch?v=pJPWi1bnl34>

-Katherine



**Brian Jay**

My dog had dog lice. Brought her to the vet and they gave her advantage. Two days later she lost control of her back legs. She was almost 16 so I thought maybe it had something to do with that as she also had some arthritis. It was so sad to see her that way. I put her down. I ask the vet if it could of had anything to do with the advantage drug and they said no. Plus she was a border collie. I feel like my dogs life was taken by this stupid drug. She was a very young 16 year old dog, still loves walks and playing outside with my son.



**Porti Dog Owner**

<https://news.wsu.edu/2013/07/08/researcher-recognized-for-drug-reaction-discovery-in-dogs/>



**Colin Lees** - It's a well known side effect, especially to dogs with the MDR1 Gene, Google MDR1



**Sarah**

I gave my parsons terrier advocate for the first time. 4 days later I found her in the middle of the night laid out on the floor unable to use her back legs and panting. I am afraid she died an hour later. She was 7 years old. I do not know if it was advocate, but I feel incredibly guilty, and would never use it again. It was prescribed, and these risks were never presented to me. Sorry to be alarmist



**jackie**

I am going through the same thing with my Collie... this product should not be used on collies or associated breeds of collies due to the MRD1 gene... this was not mentioned at all at the vets and when my dog started having problems and we as the owners started investigating this it came to our attention that kidney failure issues are a side affect and the vets did not know this... I personally think this is disgraceful as the vets should know this.. My dog is having major issues with his kidneys. we have always used Frontline with no problems so why the vet decided to change I do not know but they obviously did not do their homework and my poor dog is suffering due to their lack of knowledge



**Robert Bennett** Hi, same thing happened to mine. Beagle x lakelans terrier/jack Russel. Totally went off his legs at age of 12 months after these drops, back to normal after three days but never again. Scary what effect it had on him!



**Mrs Ingram**

I was prescribed Advocate by my vet for my boarder collie puppy when she was 4 months. She is now 6 months, and she suffered a siezure once last week and has had another one this week! She is KC registered and has no record of siezures in her blood lines. The neurological side effects from advocate, does this include siezures? After the first siezure she was taken to the vet but he had no answer to what may have caused this! Your advice would be appreciated. Thank you.



**Michelle**

Hi Mrs. Ingram,

Sorry to hear about your dog having seizures. That is a very scary event when it happens! I would advise you to contact the manufacturer, Bayer, to see if they can elaborate on the possible neurological effects that could result from using Advocate.

**Here is Bayer's contact page so you can ask them specific** questions: <https://www.bayer.com/en/contact.aspx>

I hope you get some answers so your dog doesn't have any more seizures.– Michelle



**Jenny Walsh**

Hi Mrs Ingrams I hope your dog is doing ok now. I have a thirteen year old border collie who takes advocate. But a few vets have told me that it is not generally recommended for border collies as it can cause seizures and neurological problems. My dog has been taking this for a long time so he said she mustn't carry the gene for this. I think it would be better if your dog didn't take this anymore. Take care Jenny



**Michelle**

Hi Jenny, Thank you for your information regarding using Advocate on Border Collies. That is very important information and should help other users as well.– Michellee



**Marie Kennedy Dawson**

I read that Advocate should never be given to Collie dogs!



**Kath**

Hello Where did you read this. I Have two doggies and use advocate...



**Jane**

Our cat suffered seizures immediately after advocate. She then went blind and later died of a brain tumour a year later. The chemical is a neurotoxin to brain.



**Carolyn**

we took our 8 week old Border Collie to the vet and paid for a package called "the best start" as part of this package the vet applied Advocate but did not consult with us about it she simply put it on the pups neck. One hour later he was very distressed butting his back legs and crying he was exhausted but could not sleep kept jumping about and crying. I rang the vet who said it was probably the Advocate and to wash it off which we did. He stopped being as frantic almost right away but two weeks later is still occasionally bites his legs and cries. The vet says it is nothing they have done but then they would. I am not happy I took my pup to the vet paid for care and they have possible damaged him with this product.



**Michelle**

Hi Carolyn,

Jenny just left a comment above about her experience with Advocate with her Border Collie. She says that her vet has cautioned using this product on Border Collies because it can cause seizures and neurological problems. Hopefully your dog will be fine and not have any lasting problems since you washed it off and it has had time to get out of his system.

I hope your experience and other's comments like Jenny's will raise awareness to possible dangers of this product for specific breeds like Border Collies. Thank you again for your comment.-Michelle



**Cori**

I just want to clarify it is the ivermectin in advocate that ALL herding breeds can have issues with. Like it was pointed out above they have a certain gene that makes them hypersensitive to ivermectin. If you have a herding breed (collies, shepards, heelers) you should look for heart worm prevention suggested for them. I'm not sure when this was done so I don't know if she will see this but Carolyn if they didn't tell you what was included in the puppy pack or at least have it posted then yes the vet is liable. If they did have it posted regardless of if you realized it or not they are not liable per insurance. The problem is you would have to pay a lawyer to sue them. I sure hope all the above pups are doing well. Please spread the word to other herding dog owners to have a long talk about ivermectin and their dogs.



**Michelle**

Thank you Cori for your added information. Carolyn's comment was just left a month and a half ago so hopefully she will see your comment and it will help her out. Also, I hope other herding breed owners will also become aware of the potential dangers of using certain products on their dogs.



**Katherine**

Linda, I know this post was from a long time ago, but please email me. My dog also had this type of reaction to Advantage Multi (lost use of back legs, neurological symptoms etc). I am trying to gather together those of us whose pets have had this reaction. Please contact me. I am considering legal action at this time.



**Christine**

Symptoms sound like what happened to my dog which was a stroke, an MRA didn't show why but a few days later found spleen cancer, looked like a blood clot had detached and ended up giving stroke.

Reference: <https://thedoginfoblog.com/advocate-for-dogs>

# TICK AND FLEA

## \* NEXGARD® Drug

(flea and tick prevention poison) chewable pill for dogs.

Is classified as a insecticide poison, a neurotoxin to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.... this drug is from the makers of Frontline NexGard® (afoxolaner) is available in four sizes of beef-flavored, soft chewables for oral administration to dogs and puppies according to their weight. Each chewable is formulated to provide a minimum afoxolaner dosage of 1.14 mg/lb (2.5 mg/kg). Afoxolaner has the chemical composition 1-Naphthalenecarboxamide, 4-[5-[3-chloro-5-(trifluoromethyl)-phenyl]-4, 5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-N-[2-oxo-2-[(2,2,2-trifluoroethyl)amino]ethyl.

**\*\* Symptoms of an allergic reaction to NexGard will vary from dog to dog.\*\***

**Frequently diagnosed reactions to this flea and tick treatment include:**

***Dry and flaky skin Loss of appetite Vomiting Diarrhea Lack of energy to severe neurological damages and symptoms, to death.***

## COMMENTS from consumers

~ A few hours after I gave my dog nextgard he had his first seizure and now has them a few times a week. What can I do? The vet that recommended nextgard refuses to believe that nextgard caused it since he and others that work with them use nextgard on their animals. We have taken him to Dr after first seizure and they focused on his congestive heart failure which was being and has been managed through meds for the last two years. My husband will take to vet tomorrow and demand bloodwork done for liver kidney and toxins. Plz do not use any oral flea meds. Just be patient and use topical and baths. I can see our dog isn't going to be with us much longer since nextgard was given. 1 week, 5 days ago. Shelly

~ My dog Rubyjean is getting lumps on her leg and now she is peeing on herself she can't control it I think it's from this nexgard ., 2 years ago my old dog became ill and started seizures I had to put her down because she also lost control and started peeing on herself I truly believe it's from this pill nexgard you should not sell this stuff to any dogs who are over 5 years of age older dogs can't take it I've lost my best friend and now my other dog Rubyjean looks like I May lose her to I'm heartbroken is there anything I can do to reverse treys affect 4 weeks ago Nancy S.

~ I lost my pom/poodle mix last year sept10 2017 from NexGard he was vomiting and not eating and he was having serizures something he has never had before I read that it causes serizures I called the vet and I told her he is throwing up he could be allgric to it she said just try it longer

I told them their is something not right one of the serizures lasted 45 minutes and the are suppose to have on call 24/7 ali get is answering machine and they call you back in morning is bull my baby boy is gone and I'am handid and very lost with out him he was my ears, he was my breath, he was my little helper and so, so smart and the vets and NexGard took him away. **bonnie lewis**

~ I gave my 12 year old Shih Tzu Nexguard for the first and he hasn't been the same since. He shakes and he barks incessantly unless you are with him. He faces a block wall and barks. He goes out in the yard and sits by a gate that he normally doesn't go out. I'm afraid it's killing him.

~ I believe Nexgard killed my dog. The vet pushed the sale of Nexgard on me at an appointment. I gave it to my dog and roughly a month later she died, yesterday. She hadn't been eating much a couple weeks after giving her the medication and then she had non-stop vomiting and diarrhea. Visible side-effects didn't occur right away but rather over the following weeks. 10 months ago

**Krystal W**

**Reference:** <https://wagwalking.com/condition/nexgard-allergies>

# HeartGuard Plus, HeartGuard, Nuheart

... written Dr. Jennifer Coates insecticide poison, a neurotoxin to be precise.

IVERMECTIN® Poison Product In Animals can absorb ivermectin through oral or topical exposures as well as via injection. Symptoms arise when the drug is present in the body at high enough concentrations that it crosses the blood-brain barrier and adversely affects neurologic function.

Typical signs include:

- dilated pupils
- unsteadiness when walking
- mental dullness
- drooling
- vomiting
- blindness
- tremors
- seizure
- coma

Treatment for an overdose of ivermectin is essentially symptomatic and supportive.

If the poisoning is caught early enough, decontamination is helpful (e.g., washing pets after topical exposure or inducing vomiting and/or activated charcoal administration within a few hours of ingestion). Intravenous fluid therapy, endotracheal intubation, mechanical ventilation, extensive nursing care, seizure control, application of eye lubricants if the patient cannot blink, and nutritional support may all also be necessary. In some cases, intravenous lipid emulsion therapy, which is a new but promising option for certain types of poisoning, might be worth considering.

**A pet's prognosis can be quite good if aggressive treatment is initiated in a timely manner, but because severe cases of ivermectin poisoning often require several weeks of therapy, the expense is often prohibitive ... as was the unfortunate case with my clients who chose to euthanize the last puppy in what had been their much anticipated litter.**



Dr. Jennifer Coates

# HeartGuard Plus or HeartGuard or Nuheart

**Side Effects of Ivermectin in Cats** Is classified as an insecticide poison, a neurotoxin to be precise. In cats, Ivermectin seen, side effects include:

- Agitation
- Crying
- Lack of appetite
- Dilated pupils
- Paralysis of hind legs
- Muscle tremors
- Disorientation
- Blindness
- Other neurological signs, such as head pressing or wall climbing

If your cat is receiving Ivermectin and you notice these types of symptoms, discontinue the medication and contact your veterinarian. Although you may find they will dismiss the symptoms as related to these side effects the drugs produce.

## Side Effects of Ivermectin in Dogs

In dogs, the risk of side effects associated with Ivermectin depends on the dosage, on the susceptibility of the individual dog, and on the presence of heartworm microfilaria (a larval form of the heartworm.) When used at a low dose for heartworm prevention in a dog free of heartworms, Ivermectin is relatively safe. At higher doses which may be used for treating other parasitic infections, the risk of side effects increases.

## Adverse Reactions

In clinical field trials with Heartgard Plus, vomiting or diarrhea within 24 hours of dosing was rarely observed (1.1% of administered doses).

**The following adverse reactions have been reported following the use of HEARTGARD:**

Depression/lethargy, vomiting, anorexia, diarrhea, mydriasis, ataxia, staggering, convulsions and hypersalivation, Dilated pupils, Muscle tremors, Blindness, Incoordination, Lethargy, Lack of appetite, Dehydration (from fever).

\* Certain dogs breeds are more sensitive to this poison and have showed adverse reactions which included mydriasis, depression, ataxia, tremors, drooling, paresis, recumbency, excitability, stupor, coma and death.

# FRONTLINE - Fipronil (Frontline gold Frontline Plus )

Fipronil, a chemical insecticide, is the primary active ingredient in the various Frontline products.

**Is classified as a insecticide poison, a neurotoxin to be precise.**

## Frontline Side Effects for DOG

**Its active ingredient, fipronil, lists other potential concerns when used on canines.**

- **Skin problems** – As stated earlier, fipronil can cause irritation to your dog's skin. This irritation can go beyond simple itching as the scratching can cause ulceration and open sores.
- **Nervous system damage** – Fipronil is a neurotoxin, causing damage to the fleas' nervous system. It can have the same effect on your dog, causing symptoms such as convulsions, body twitches, loss of appetite, unsteady gait and other effects.
- **Carcinogen** – Fipronil has been shown to cause **thyroid cancer in dogs**. Because of its carcinogenic qualities, it is important for the person applying the Frontline product to thoroughly wash their hands with soap and warm water to adequately remove the product from their skin.
- **Organ damage** – Autopsies have shown an accumulation of fipronil in canine **livers and kidneys DAMAGES**, causing an increase in organ weight. Fipronil has also been shown to alter the levels of hormone secreted by the thyroid gland **CAUSING HypoThyroid conditions**.
- **Infertility** – Fipronil has also been shown to have an effect on a dog's fertility. Its use has led to smaller litter size, difficulty in conception, decreased litter weights and **increase in litter deaths**.

Frontline Side Effects For CAT *Since the active ingredient fipronil works as a pesticide, it's important to avoid using the treatment along with other pesticides. Pets may experience redness or irritation at the site of application. Common side effects include allergic reactions to the product, in pets hypersensitive to the ingredients. If the cat develops rash, swelling or hives associated with allergic reactions, it's important to seek prompt vet care. Any adverse reactions or side effects using Frontline are usually due to incorrect administration or dosage. However, all pets are individuals and may be hypersensitive or allergic to the ingredients. If you notice any of the following side effects take your pet to the vet immediately:*

- Swelling particularly of the face, lips and tongue
- Rash or hives
- Lethargy
- Vomiting or severe/lengthy diarrhoea
- Excessive salivation
- Uncoordinated movements
- Seizure
- Unresponsiveness or coma
- Overdose may cause certain reactions and pets require emergency care
- The product is not safe for use in sick or debilitated pets

## Pet owners COMMENT after using these products

**Lorene** 12 Aug 2017

- I use to use Frontline Plus on my cat, but yesterday vet gave me frontline gold. Anyway was in the vet office for 2 hours this morning, as my usually noisy up at 5 AM cat, was hiding under the bed this morning and had vomited during the night. After almost \$400 of fluids, anti nausea, and blood panel, she is still not meowing, eating or moving around much. Not applying this stuff ever again.

**Jim** 12 Aug 2017

- Mt cat almost died 24 hours after the first application of Frontline Gold. It has some additional chemicals in it that Frontline Plus does not that are extremely toxic. After numerous vet visits and a week or so, the cat pulled thru. He is back to Frontline Plus with no problems.

**Jo** 23 Aug 2017

- I gave my 20 year old cat Frontline after reading the instructions and telling my vet I was doing it. On both occasions he has gone off his food and had diarrhoea which ARE on the list of side effects, but he goes blind for 24 hours about 48 hours after application... this was certainly NOT on the list. It's very scary and distressing but does not last .... be warned everyone!!

**David** 22 Jan 2018

- I applied Frontline on my dog 13lbs a month ago. I have had him to two different vet's in last two weeks. They can't find anything causing his situation most all the Symptoms as everyone else. Frustrated and it was time to reapply the Frontline. It dawned on me to look up and find what you all are talking about. Being real sure this is what is going on with my dog. Called the vet. And they say doubtful. I don't understand why the vets dont stand up for it. I know it is the problem. All the Symptoms reappear. Strange. Battery eyes, shaking, loss of appetite, lethargic, ect. I need to find a vet willing to accept the diagnosis and somehow treat.

**Eleanor** 24 May 2018

- Sad stories indeed. I've had the same experience with **HeartGuard** for heart worms.

My little dog began to have seizure-like symptoms which worsened over the months. Finally after the last dose I gave her she couldn't even stand within two hours. The vet never mentioned it might be the ivermectin. He still refuses to believe that was the cause. After I finally researched the product I found out the truth. All of these products contain neuro toxins that are supposed to kill the parasites and not cross the blood brain barrier of the animal. But with some dogs and especially sickly or old ones, it does! My dog had what they called ataxia. Staggering, shaking, seizure-like symptoms, eyes flitting back and forth rapidly, until finally after the last time we thought we'd have to put her to sleep. But we didn't, and it took 6 weeks in a dog baby buggy and hand feeding because she was blind and couldn't hold her jerking head still enough to eat on her own. It ends well though. I quit giving it to her and now I use Wondercide. The one I use has cedar oil and hydrated silica, which is like diatomaceous earth. It kills the fleas! Really. Has to be applied quite often if it's to repel mosquitos too, but in winter a couple of times a week handles fleas. The fleas just sort of dissolve. The little dog recovered very well after that and loved having the cedar smell on her. Not all of them like it that much but the point is, it will do no harm and it works if you're willing to go to the trouble. I never want to poison my sweet dogs again, so I do.

- ..... Scroll down to read the many more COMMENTS of side effects from people using these products (link below)

**Reference:** <https://www.petbucket.com/blog/62916/the-side-effects-of-frontline-flea-tick-prevention.html>

# TICK, FLEA AND WORMER

## TRIFEXIS<sup>®</sup> (spinosad + milbemycin oxime)

\* Used as a Wormer, Flea, Heartworms –

**Is classified as a insecticide poison, a neurotoxin to be precise.**

**A Neurotoxin to be precise.** These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function. It is the monthly, beef-flavored tablet that poisons your pets blood so it will also be that toxic in the system it will also kills fleas and prevents **heartworm** disease, and treats and adult hookworm\*, roundworm and whipworm infection.

### WHAT ARE THE SIDE EFFECTS OF TRIFEXIS?

The most common adverse reactions based on post-approval adverse drug event reporting are .. *vomiting, lethargy, itching, loss of appetite, diarrhea, trembling / shaking, incoordination, seizures, excessive salivation and redness of the skin.*

## NEWS FLASH

According to FDA records, pet owners have reported 700 dog deaths.

That's 120 more than the deaths linked to Chinese chicken jerky treats. Like the treats, Trifexis is suspected. [www.wsbtv.com](http://www.wsbtv.com) > news > local > o...

### Owners blame 700 dog deaths on Trifexis - WSBTV.com

The most frequently reported adverse reaction in dogs in the TRIFEXIS group was vomiting. In the US field study, one dog administered TRIFEXIS experienced a single mild seizure 2 1/2 hours after receiving the second monthly dose ...Oct 21, 2020

[www.drugs.com](http://www.drugs.com) > vet > trifexis

Trifexis for Animal Use - Drugs.com

## SIMPATICO TRIO<sup>®</sup> Chews

**Is classified as a insecticide poison, a neurotoxin to be precise.**

**Contain insecticide poison -**

200mg pyrantel (embonate), 48mg sarolaner, 0.96 mg moxidectin.

**Sold and advertised as – flea, tick, heartworm and wormer.**

### Side effect Warnings

Most common adverse reactions being vomiting and diarrhea—both symptoms causing stomach pain digestive tract reactions to toxins, rather than his nervous system.

Dogs have also had more serious reactions such as *tremors, muscle twitching, and seizures.*

### **Let's look at "Heartworm" - A realistic Look**

**Heartworm is of concern to all dog owners. It is not, however, the unavoidable scourge that many vets and pharmaceutical companies would have us believe.**

Most people do not have any idea what is involved in a dog getting infected with the Heartworm larvae. This article should educate you on the life cycle of this larvae and ways to prevent any effects if your dog is infected. If your dog is reasonably healthy, eating a proper real food diet, is over use "vaccine free", is not currently on any Heartworm "preventives" or monthly chemical flea and tick preventives, their immune system should be strong enough to fight off a heartworm infestation with none of the larvae reaching adulthood.

In the event that some of the larvae should manage to get past your dog's immune system defenses and survive to adulthood, it is still far from a death sentence. They will be much weakened, and the truly healthy dog will make a less hospitable host. Even at the adult stage, a healthy dog, possibly with some veterinary assistance, should be able to fight them off with no lasting ill effects to their health.

**Nor are the currently available medical cures (arsenic drugs) for Heartworm more dangerous than the monthly "preventives," as the veterinary and pharmaceutical industries would have us believe.**

**Mosquitoes are the only known carrier of the infective stage of the heartworm larvae.** In some parts of the country mosquitoes are a problem all year round, while in others this problem is limited to the warm months. There are many sites out there on the internet with lots of "information" on heartworm.

The problem with much of it is that it is either wrong or incomplete. **Much of it is intended to mislead and frighten you.** One site says simply that a mosquito bites an infected dog then bites another dog and infects it.

**While that, to some extent, in the most general of ways, is true, it is very misleading because the entire process is much more complicated and time consuming than this "explanation " would lead one to believe.**

Another explanation says that the mosquito injects the heartworm into your dog. This is simply false. Long or semi haired dogs, or dogs that sleep and live inside a house do not need to on heartworm drugs, but you can safely use alternatives if so desired.

## \* BRAVECTO® chewable tablets

\* Given as a poison to kill fleas, poisons ticks, and mites - demodex, sarcoptes and ear mites) which is classified as an insecticide poison.

A Neurotoxin to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.

### Comfortis and Capstar drugs

Bravecto is a relatively new product released in 2014 which is used for **fleas and ticks**. It contains the active ingredient fluralaner, a type of isoxazoline which maintains protection for 12 weeks, meaning you only have to use the medicine as a preventative roughly once every 3 months. That means it has a high amount of pesticide chemical toxins in it.

## How does it compare to Comfortis and Capstar?

Comfortis and Capstar are two other popular flea treatments. Unlike Bravecto, Comfortis does not kill ticks and needs to be given every month, and Capstar is not effective as a flea preventative. Which medicine you choose will come down to personal preference and how well your dog tolerates each of them. As you may have read online, many owners have seen Bravecto has severely harmed their dog. **Research before giving your dogs or cats edible pesticides, once eaten there is NO ANTIDOTE.**



These are the most recent numbers of reported side effects and deaths after treatment with the tick and flea treatments **Bravecto, Nexgard and Simparica**. Please note that only about 1 % of the side effects is actually reported by veterinarians and dog / cat owners.

Most vets will simply deny that the symptoms are related to administration of these pesticides  
BROADCAST FROM THE NETHERLANDS : THE ANTI BRAVECTO PETITION WITH 40.000 SIGNERS HAS  
BEEN DELIVERED AT THE MINISTRY - please don't forget to share, to spread the word...

**The petition will continue, so you can still sign it via [www.bravectopetition.org](http://www.bravectopetition.org)**

## Some of CONSUMERS COMMENTS

**Bravecto Insecticide poison (to prevent or kill fleas, but is killing pets too**

Will never use **Bravecto** again. June 23, 2017

I gave my 2 dogs **Bravecto** after seeing an advert on media saying it lasted for 3 months. I thought perfect because I always forget to get them their tablets at the end of the month. Within hours of my 3yo Maltese Shih Tzu and 1yo Maltese poodle having the 1 tablet each they were struggling to breathe and were shaking and panting erratically. They also both were very lethargic and wouldn't eat. I was very worried and looked online for any information and then called the vet after reading all the terrible reviews on this option for Flea and Tick Control. They advised they had no idea about any adverse reactions from this product and told me to just keep an eye on them. I wish I had reviewed prior to giving my dogs this tablet. In the future, I will check reviews prior to purchasing. It took over a week for the shaking to subside and their breathing came well after about 3-4 days. After those reactions, I would not recommend this product to anyone. I will never use Bravecto again for my pets.

**POISON** June 07, 2017

I have never written a review of anything but I must warn everyone....my beautiful dog Toby died Sunday, May 29th, 2017 from me giving him this POISON on a recommendation from my vet. He was a perfectly healthy, lovable and perfect lab mix (just examined fully) a few days after giving him the drug he stopped eating and began to vomit, I rushed him to the vet they did x rays, and a pancreatic scan and both were good, we went home and less than two days later he couldn't move and had become completely delirious so we rushed him to the animal hospital where his liver values were found to be through the roof, they started iv meds but unfortunately it was not enough to save our boy and he passed overnight. PLEASE SAVE YOUR DOGS LIFE AND DO NOT GIVE THIS MEDICATION. NO DOG SHOULD HAVE TO GO THROUGH WHAT MY DOGGY DID AND NO FAMILY SHOULD HAVE TO SEE THEIR DOG GO THROUGH IT.

**May Work For Some, But Deadly For Others** May 17, 2017

Never again will I give this product to any of my 3 dogs. Sure, they don't have fleas or ticks, but after taking it, it made one dog sluggish for days, and another almost died. He went from being a perfectly healthy dog to lethargic, depressed and he completely stopped eating within weeks of administering Bravecto. His abdomen filled with fluid, he became jaundiced, developed mouth sores, his nose started dripping blood fluid...all of this out of nowhere. He is a bag of bones, from not having an appetite. The Vet ran lab work and his liver enzymes were off the charts. Several rounds of Xrays revealed his liver was extremely enlarged.

... The only thing that had changed was I administered Bravecto, based on recommendations by my dog's veterinary office. This was the first and ONLY time I will ever use this product. Over \$1000 later, lots of TLC and hand feeding, a whole new home-cooked diet, and expensive daily medications that were never needed before, and it's still a touch and go roller coaster ride, but we are nearing the end of the 3 months and he's still here. Just remember, this medication lasts for THREE months, so if it doesn't agree with your dog, it's not going anywhere for awhile, and there are no antidotes if it affects your pet. Please think long and hard before giving this to your furbaby!

### [I need help HAMPL](#) 25th June 2017

Please include a liver detox and magnesium glycinate for my 3 yr old BostTerrierier has suffered 2 **seizures** since being administered "**Bravecto flea and tick**", vet believes unrelated, but the dog has never had seizure prior to this parasite preventative. many many thanks.

**Emma** (Melbourne)

## NATURAL TREATMENT FOR DAMAGES FROM USING Bravecto

### **"BRAVECTO"** damages causing neurological and organ damages.

Your dog or cat will most likely have all of the below damages. As we have been helping so many severely damaged pets from this drug, some are dying before we could help them.

#### ~ **Vitamin B12 - deficiency anemia.**

Anemia caused by low intrinsic factor production is called "**Pernicious Anemia**".

Add **vitamin B12** in meals twice daily. use the Liquid B12 (from VIMERGY.com) Give 1 to 2 drops in the gum of pets mouths, three to four times a day until improvement is seen that can reduce to 1 drop twice a day for a month. \* Plus **Vitamin C** (Liposomal gel or liquid)

#### ~ **Pancreatitis (acute inflammation)** \* *this is very painful*

Use [HAMPL Acute Pancreatitis 72-2 30ml\(1oz\)](#)

#### ~ **Chemical Poisoning Detox**

Use [Chem Detox 110 30ml](#) with Activated Charcoal powder

#### ~ **Neurological Brain Damages** ( as this toxic chemical is neurotoxin)

Symptoms-CNS (central nervous system)Cerebral Ataxia,e.g.paralysis, head nodding, paralysis agitates- tremor, impaired muscular coordination.

1. [HAMPL Brain Nerve 94-1 30ml drops](#)

2. [HAMPL Paralysis Gone 30ml drops](#)

3. [HAMPL Systemic Infection 94-3 50ml liquid](#)

~ **Relax muscles spasms and shakes.**

Magnesium (mineral) e.g ReMag Magnesium drops or Magnesium Glycinate powder.

\* See pdf HELP SHEET beside the AN284 ReMag Magnesium product

~ **Diarrhea** - Use slippery elm powder diluted and mixed in some liquid and oral syringe.

Product [AN118 Slippery Elm](#) or buy from Health food stores.

\* See pdf HELP SHEET found at [AN118 slippery product for instructions.](#)

~ **Organ Failure** \* *if already have a weak heart or kidney function*

Use [HAMPL Acute Nephritis 19-2 30ml](#) \* *Acute Inflamed Kidneys*

Use [HAMPL Congestive Heart Failure 22-1 50ml](#) \* *Heart function*

~ **Seizures** ( or jerking, spasms, tremors)Use [Canine Seizure 026-1 drops](#) and **Magnesium** and B6\* See [pdf HELP SHEET AN026 Seizures](#) for further support and healing.

Needing more help and support, please contact your practitioner.

## SENTINEL tablets

### Is classified as a insecticide poison, a neurotoxin to be precise.

A Neurotoxin to be precise. These toxins are poisonous or destructive to nerve tissue. Neurotoxins are an extensive class of exogenous chemical neurological insults that can adversely affect function.

**Drug Name:** Sentinel

- **Common Name:** Sentinel®
- **Drug Type:** Parasiticide
- **Used For:** Used for fleas, ticks, heartworms, mites
- **Species:** Dogs, Cats
- **Administered:** Tablets

**Sentinel® contains ~ Milbemycin Oxime**, which is effective against internal parasites such as roundworms, hookworms, whipworms, and young heartworms. It works by interfering with the **central nervous system** of the heartworm larvae. It is not effective against any stage of the flea life cycle. ~ **Lufenuron**, which prevents egg production in fleas. Lufenuron does this by preventing the flea larva from being able to make chitin, which makes them unable to produce an exoskeleton and grow. Sentinel is not effective against adult fleas, but prevents growth of the infestation on your pet. Sentinel® heartworm, common intestinal worms and fleas. As is true with any prescription medication,

### Sentinel does have serious side effects.

**The following side effects have been observed in dogs taking Sentinel.**

- Diarrhea.
- Depression, lethargy, or weakness
- Drooling.
- Loss of appetite.
- Dilate (big) pupils
- Vomiting.
- Staggering, stumbling or collapse.
- Seizures or coma

## **Gastrointestinal Side Effects**

Side effects after taking Sentinel for dogs are extremely rare. However, of those dogs that do experience adverse reactions, diarrhea, excessive drool, loss of appetite and vomiting are common. Let the dog rest and avoid strenuous activity, especially following a meal, because activity can increase the possibility of regurgitation. Make sure he's drinking to avoid dehydration.

## **Skin Reactions Following Sentinel for Dogs**

Though Sentinel is an oral medication, it may cause itchy skin, oily skin and hives. Shampooing your dog with an oatmeal-based pet-safe shampoo can help relieve the irritation. If the condition continues, contact your veterinarian.

## **Signs of an Allergic Reaction to Sentinel**

If your dog is having an allergic reaction to at least one of the active ingredients in Sentinel, symptoms include:

- Cold extremities
- Coma
- Facial swelling
- Pale gums
- Severe vomiting
- Sudden, explosive diarrhea

## PROFENDER®

Cat - is insecticide poison used for hookworm, roundworm, and tapeworm

**Is classified as a insecticide poison, a neurotoxin to be precise.**

**CLINICAL STUDIES** - Using Topical Solution to cats.

**Adverse reactions** reported by the cat owners included licking/excessive grooming in 18 cats (3.0%), scratching treatment site in 15 cats (2.5%), salivation in 10 cats (1.7%), lethargy in 10 cats (1.7%), alopecia in 8 cats (1.3%), agitation/nervousness in 7 cats (1.2%), vomiting in 6 cats (1.0%), diarrhea in 3 cats (0.5%), eye irritation in 3 cats (0.5%), respiratory irritation in 1 cat (0.2%) and shaking / tremors in 1 cat (0.2%). All adverse reactions were self-limiting.

### **Laboratory effectiveness studies:**

One cat died 10 days after receiving PROFENDER Topical Solution. The necropsy showed chronic active cholangiohepatitis (is a word that describes inflammation of the liver and bile ducts). (See PRECAUTIONS).

Other cats treated with a vehicle placebo (formulation minus the active ingredients) showed salivation, gagging, lethargy and a swollen tongue

## REVEOLUTION Plus®

Selamectin and Sarolaner **classified as a insecticide poison and neurotoxin**

**(affects the nervous system-seizures and in coordination etc).**

### **Revolution side effects:**

**Cat and dogs experienced digestive upset as a result of using this product.**

*Vomiting, loose stools, diarrhea with or without blood, anorexia, lethargy, salivation, tachypnea, and muscle tremors, **in coordination, fur damage and loss on site of poison was applied, are other Revolution side effects listed in packet.***

**CAT** Side effects of Revolution Plus ...**lethargy, diarrhea, emesis, eczema, skin lesions, anorexia, pruritus, conjunctivitis, sneezing, administration site hair changes, or administration site lesions. tremors, ataxia, and seizures. Reactions have occurred in cats with or without a history of neurologic disorders.** As sarolaner chemical poison is one of the ingredients in Revolution Plus is a member of the isoxazoline class. **This class has been associated with neurologic adverse SIDE EFFECTS IT IS A NEUROTOXIN to the brain and nervous system) ... reactions including tremors, ataxia, and seizures.** Neurologic adverse reactions have been reported in cats receiving isoxazoline class drugs, even in cats without a history of neurologic disorders.

# TICK FLEA COLLARS & MANGE MITE

**Poison products** - Demodex mites or skin demodex There are ALWAYS alternatives that are safe - TICK AND FLEA e.g <https://keepthetailwagging.com/flea-and-tick-collars/>

**Is classified as a insecticide poison, a neurotoxin to be precise.**

## AMITRAZ® - Insecticides Poisoning in Dogs

**Brand Names and Other Names of Amitra**

Mitaban® (Upjohn),

Preventic® (Allerderm/Virbac),

Tactic® (Hoechst/Roussel) and various generic brands.

### Uses of Amitraz for Dogs

- Amitraz is primarily used to treat the skin disease demodectic mange.
- It is also used for tick prevention.

**Mechanism of action of Amitraz poison is:** The acaricidal and insecticidal activity effect on **octopamine receptors** of the nerve cells in the brain. Parasites become hyperexcited, paralyzed and eventually die. This too can cause central nervous system paralysis in your pet.

**Symptoms of Amitraz insecticide poisoning (Amitraz toxicosis) in dogs include:**

Low blood pressure.

Low body temperature.

**Elevated** levels of blood glucose.

Ataxia.

Decreased heart rate.

**Vomiting.**

Diarrhea.

Seizures.

**The most common symptoms seen in a dog include:**

Vomiting

Diarrhea

Lethargy

Weakness

Staggering

Disorientation

Hypothermia

Abdominal pain

Light or severe depression

**Reference:** <https://wagwalking.com/condition/amitraz-insecticides-poisoning>

## SERESTO® Collars

Active ingredients, 4.5% Flumethrin and 10% Imidacloprid classified as a insecticide poison and neurotoxin (*affects the nervous system-seizures and in coordination etc*). According to the NPIC, all kinds of pesticides can cause health problems in your pet. This includes products such as shampoos, sprays, and even pills.

**Here are the side effects associated with Seresto collars and other pesticide products:**

Rash

Vomiting

Diarrhea

Trouble walking|

Tremors and seizures

Excessive fatigue

Excessive drooling

Fatal

Some animals may be more sensitive to certain pesticides than others. Thus, always consult with your vet if you notice any health changes in your pet.

**APOQUEL® Drug** prescribed mainly for skin dermatitis issues for Canines. (atopic)

**Pruritus control (not healing) associated with allergic dermatitis of atopic dermatitis in at least 12 months of age.** \* too dangerous to give to young pets or felines.

### Adverse Reactions Studies

In a masked field study to assess the effectiveness and safety of oclacitinib for the control of atopic dermatitis in dogs, 152 dogs treated with APOQUEL and 147 dogs treated with placebo (vehicle control) were evaluated for safety. The majority of dogs in the placebo group withdrew from the 112-day study by Day 16. Adverse reactions reported (and percent of dogs affected) during Days 0-16 included diarrhea (4.6% APOQUEL, 3.4% placebo), vomiting (3.9% APOQUEL, 4.1% placebo), anorexia (2.6% APOQUEL, 0% placebo), new cutaneous or subcutaneous lump (2.6% APOQUEL, 2.7% placebo), and lethargy (2.0% APOQUEL, 1.4% placebo).

In most cases, diarrhea, vomiting, anorexia, and lethargy spontaneously resolved with continued dosing. Dogs on APOQUEL had decreased leukocytes (neutrophil, eosinophil, and monocyte counts) and serum globulin, and increased cholesterol and lipase compared to the placebo group but group means remained within the normal range. Mean lymphocyte counts were transiently increased at Day 14 in the APOQUEL group. Dogs that withdrew from the masked field study could enter an unmasked study where all dogs received APOQUEL. Between the masked and unmasked study, 283 dogs received at least one dose of APOQUEL. Of these 283 dogs, two dogs were withdrawn from study due to suspected treatment-related adverse reactions: one dog that had an intense flare-up of dermatitis and severe secondary pyoderma after 19 days of APOQUEL administration, and one dog that developed generalized demodicosis after 28 days of APOQUEL administration.

Two other dogs on APOQUEL were withdrawn from study due to suspected or confirmed malignant neoplasia and subsequently euthanized, including one dog that developed signs associated with a heart base mass after 21 days of APOQUEL administration, and one dog that developed a Grade III mast cell tumor after 60 days of APOQUEL administration. One of the 147 dogs in the placebo group developed a Grade I mast cell tumor and was withdrawn from the masked study. Additional dogs receiving APOQUEL were hospitalized for diagnosis and treatment of pneumonia (one dog), transient bloody vomiting and stool (one dog), and cystitis with urolithiasis (one dog). In the 283 dogs that received APOQUEL, the following additional clinical signs were reported after beginning APOQUEL: pyoderma (12.0%), non-specified dermal lumps (12.0%), otitis (9.9%), vomiting (9.2%), diarrhea (6.0%), histiocytoma (3.9%), cystitis (3.5%), anorexia (3.2%), lethargy (2.8%), yeast skin infections (2.5%), pododermatitis (2.5%), lipoma (2.1%), polydipsia (1.4%), lymphadenopathy (1.1%), nausea (1.1%), increased appetite (1.1%), aggression (1.1%), and weight loss (0.7%).

## Side Effects of Apoquel for Dogs

(as its not recommended for cats do to acute toxicities reactions are fatal)

Just like any other drug, Apoquel comes with its own set of side effects and possible reactions. Because of its immunosuppressive properties, it can increase your dog's vulnerability to infection. This includes demodicosis infections and neoplastic health conditions.

### Other potential side effects of a pet taking Apoquel include:

- Vomiting
- Diarrhea
- Appetite loss
- Lethargy
- New cutaneous or subcutaneous lumps
- Decreased leukocytes
- Decreased globulins
- An increase of lipase and cholesterol

### Also experienced:

- Bloodied diarrhea
- Pneumonia
- UTIs
- Skin and ear infections
- Histiocytomas
- An increase in appetite
- An increase in aggression
- And developed polydipsia

These side effects are similar to what a dog may experience when taking steroids. Note that some of the side effects from taking this drug may resolve on their own with continued use, as the body adjusts.

Oclacitinib (Apoquel™) **used for:**

- Control of atopic dermatitis without cortisone side effects
- Animals at risk of diabetes, cushings disease, obesity, pancreatitis
- Animals prone to infections
- Dogs only **No Cats (not suitable)**

## COMMENTS from consumers

.... My dog was on Apoquel for one week when he began having seizures. We stopped the Apoquel, treated the seizures with phenobarbital for 5 weeks, and he was seizure free. He began chewing himself again, the dr said it was safe to put him back on Apoquel, which my dog did another round of for a week. **Skye Leibee**

... He had a seizure, out of the blue, in the early evening of February 18. He subsequently had 2 more, within 2 hours of the first one. The last seizure came at about 1:30 am, on the 19th, which lasted about half an hour by the time we got him to the emergency vet. I had to say goodbye to my spunky dog of 12 years, who suffered a horrible grand mal seizure. Now I know he was an "older" dog, but he was otherwise healthy, with no history of seizures.

**Reference:** <https://dogshealthproblems.com/apoquel-side-effects-in-dogs/>

# TOPICAL SKIN TREATMENT

## antifungal, antibacterial shampoo

### \* MALASEB® - Topical Skin antibacterial and antifungal

( chemical Miconazole Nitrate 2%, Chlorhexidine Gluconate 2% POSION)

#### **Warning of side effects.**

##### **Skin Irritation (*allergy reaction*)**

When treated long-term or in dogs with very sensitive skin, irritation may develop. Watch your dog's skin for signs of further irritation directly related to the product. Take note of the skin's surface before the medication or shampoo is applied and notice if the skin deteriorates after use. Look for additional redness, scratching, and oozing. Malaseb should never be directly applied to open wounds.

##### **Doing so may result in further damaged skin cells and infection.**

##### **Nausea and Vomiting (*gastro intestinal pain*)**

Malaseb should not be ingested. Careful attention should be paid to keep your dog from licking the areas where the malaseb was applied until the product has dried. Typically, if you are using a shampoo, once the fur is completely dried, your dog can run around unsupervised. If you use a spray or towelette, the area that is wet must be thoroughly dry before your dog can lick that spot. Distract your dog with a game of catch or take it on a long walk, ideally in the sun, to aid drying. If ingestion occurs, consult your veterinarian.

##### **Eye Inflammation**

Malaseb is not safe for a dog's eyes. When using a spray, hold your dog's head to the side or have another person help you hold your dog to avoid getting malaseb in the eyes. If, despite your efforts, some spray does enter your pup's eyes, wash out immediately with a cold water. Long term eye damage is rare but precautions should be taken to avoid unnecessary pain and discomfort. If eye redness develops or if the affected area is close to your dog's eyes, consult a veterinarian.

# ANTIBIOTICS drugs, creams etc

## ANTIROBE® Clindamycin Antibiotic drug

(brand name is called clindamycin) **Oral, Topical, Injectable**

Clindamycin is an antibiotic and treats infection. This medication is available as a topical cream, topical gel, topical solution, topical foam, topical swab (pledget), vaginal cream, vaginal suppository, oral capsule, oral liquid, and granules for oral solution. This medication is also available in injectable forms to be given directly into a vein (IV) or a muscle (IM) by a healthcare professional.

Common side effects of clindamycin ... include nausea, vomiting, loss of appetite, diarrhea, joint pain, heartburn, pain when swallowing. It's also possible that your pet might have a more

extreme allergic reaction to the medicine. Serious side effects have been reported with clindamycin. See the "Clindamycin Precautions" section. Because clindamycin therapy has been associated with severe colitis which may end fatally, it should be reserved for serious infections where less toxic antimicrobial agents are inappropriate. This is not a complete list of clindamycin side effects. Ask your doctor or pharmacist for more information. Tell your doctor or Vet if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

### Antirobe Precautions-

#### **Antirobe Oral DOSAGE:**

Serious side effects have been reported with oral clindamycin including the following:

- Abdominal pain
- Gastrointestinal upset
- Pain or difficulty when swallowing
- White patches in the mouth
- Blisters
- Rash
- Hives
- Yellowing of the skin or eyes
- Decreased urination

#### **Antirobe Topical:**

Serious side effects have been reported with topical clindamycin including the following:

- Dry, itching,  
burning, or peeling skin
- Headache
- Diarrhea
- Abdominal pain or gastrointestinal upset
- Thick, white discharge or burning, itching, or swelling of the vagina with use of vaginal clindamycin products.

## Antirobe Injectable:

Serious side effects have been reported with injectable clindamycin including the following:

- Abdominal pain
- Gastrointestinal upset
- Pain or difficulty when swallowing
- White patches in the mouth
- Blisters
- Rash
- Hives
- Yellowing of the skin or eyes
- Decreased urination

### **Do not take clindamycin if you or your pet:**

are possibly having an allergic reaction to clindamycin or to any of its ingredients  
are allergic to lincomycin (Lincocin)

## VERAFLOX® pradofloxacin Antibiotic drug

Oral Suspension for Cats is the next step in veterinary antibiotic therapy for cats. It is approved for the **treatment of skin infections** (wounds and abscesses) in cats caused by susceptible strains of **Serious Side Effects** Veraflox is for cats only. Veraflox can cause **serious side effects if given to dogs.**

Side Effects of Veraflox can include : *diarrhea, leukocytosis with neutrophilia (high white blood count), elevated CPK levels, sneezing, hematuria (blood in urine), hypersalivation, pruritus (severe itching of the skin), lack of appetite, lethargy, cardiac murmur, reclusive behavior, vomiting, bacteriuria (bacteria in urine), polydipsia (great thirst), and upper respiratory infection.*

## \* METRONIDAZOLE® Antibiotic drug

Metronidazole is a prescription antibiotic used in dogs and cats to treat various conditions such as **inflammatory bowel disease, nonspecific diarrhea disorders, infections caused by Giardia, and periodontal disease.** SIDE EFFECTS - **The more common side effects that can occur include:** headache, loss of appetite, nausea or vomiting, diarrhea, heartburn, cramping in your stomach area, constipation, metallic taste.

## \* DOXYCYCLINE®

*A Broad spectrum antibiotic (tablets or oral liquid )ALSO CALLED (Vibramycin®, Doxychel®, Doxy Caps®, Bio-tab®, Monodox®, Doryx®, Doxirobe®)*

*Doxycycline is a broad spectrum antibiotic that is a member of the tetracycline family. It is used to fight bacterial infections in dogs and cats. Doxycycline is used to treat many different bacterial infections including, leptospirosis, toxoplasmosis, mycoplasma, psittacosis, and tick borne diseases including Lyme disease, ehrlichiosis, and Rocky Mountain spotted fever. Treatments using Doxycycline are generally termed as a long term treatment.*

### **SIDE EFFECTS and Drug Reactions***Doxycycline may result in these side effects:*

*Nausea, Vomiting, Swallowing difficulties, If Doxycycline makes your pet nauseous, try administering it with food. If it makes your pet have breathing or swallowing problems, follow tablet with a small amount of water. Doxycycline may react with these drugs: Antacids, Anticoagulants, Bactericidals, Barbiturates, Cathartics, Gastro-intestinal protectants, Aminophylline, Digoxin, Insulin, Iron dextran, Kaolin/Pectin, Methoxyflurane, Sodium bicarbonate, Theophylline .*

**USE CAUTION WHEN ADMINISTERING THIS DRUG TO YOUNG PETS, PREGNANT OR LACTATING PETS, OR PETS WITH SEVERE LIVER DAMAGE. (AS THIS DRUG DAMAGES THE LIVER EVEN MORE SO)**

## \* CLAVAMOX® Amoxy Clavulanic acid

**Broad spectrum antibiotic for cats and dogs.**

Clavamox is a mix of amoxicillin and clavulanic acid, two medications that work in combination to fight bacterial infections. Clavamox may be used to treat skin wounds and abscesses, urinary tract infections, tooth infections, pneumonia, injuries caused by bites and other infections.

The medication cannot be used to treat infections caused by a virus or parasite

### **Side effects are commonly seen:**

- Diarrhea or loose stool
- Allergic reaction (labored breathing, hives, etc)
- Loss of appetite.
- Vomiting/Nausea.
- Gastrointestinal pain and damages.
- Liver damages - elevations in serum transaminases (ALT and/or AST). Hepatic dysfunction (including cholestatic jaundice and hepatitis, increases in ALT and/or AST, serum bilirubin, and/or alkaline phosphatase.

### **And many others**

<https://www.drugs.com/sfx/amoxicillin-clavulanate-side-effects.html>

## CONVENIA<sup>®</sup> drug

Convenia is a veterinarian-prescribed antibiotic medication indicated to treat certain types of bacterial infections in cats and dogs. It is administered as one subcutaneous injection beneath the surface of the skin, and is effective for up to 14 days, according to Pfizer Animal Health, the pharmaceutical distributor of this medication. Cats and dogs must be at least 4 months old before receiving Convenia as its side effects are dangers.

### Side Effects of Convenia

**Nausea or Vomiting** - After receiving Convenia, cats and dogs may develop upset stomach side effects, including nausea or vomiting. During clinical trials for this medication, vomiting was observed in approximately 5 percent of dogs and 8 percent of cats, reports Drugs.com. Upset stomach side effects may also contribute to a temporary decrease in appetite. Pet owners may notice that their pet is uninterested in eating at mealtime or won't eat favorite treats. These side effects typically subside within a few days of treatment.

**Diarrhea** - Convenia may cause diarrhea side effects, PetPlace.com reports. Pet owners may notice that they need to clean the litter box or let their dog out more frequently than usual. Depending on the severity of diarrhea side effects, certain pets may accidentally soil furniture or flooring due to sudden bowel movement urges. If the diarrhea persists or becomes severe, or if pet owners notice blood in their pet's stools, the affected cat or dog should be taken to a veterinarian as soon as possible.

**Fatigue or Lethargy** - In clinical trials for this medication, lethargy was observed in approximately 4 percent of treated cats and dogs, according to Drugs.com. Affected pets may seem tired or may not be as playful as usual. Certain cats treated with Convenia may display unusually hyperactive behavior, PetPlace.com warns. Such side effects generally resolve within a few days following initial treatment with Convenia.

## \* BAYTRIL<sup>®</sup> (Enrofloxacin)

broad spectrum antibiotic

**Side effects are commonly seen:**

Loss of appetite.

Vomiting.

Diarrhea.

Lethargy.

Convulsions.

Seizures in pets with CNS disorders.

Cataracts if given long-term.

We have a safe and very fast acting natural medicines to treat infections

**HAMPL Infection Fighter 71 50MI**

# FUNGAL TREATMENT DRUG.

## FLUCONAZOLE® Drug

**is an antifungal drug effective against fungus and yeast (e.g Ringworm treatment).** It is typically used for skin infections, fungus in the nailbeds, and more intense fungal infections including Blastomycosis and Histoplasmosis. It is related to Ketoconazole, but is able to more effectively cross the brain-blood barrier, making it more successful in treating fungal infections of the central nervous system. Fluconazole is also USED against ringworm, but is typically reserved for more serious infections.

### Side Effects and Drug Reactions

Fluconazole has been found to be very safe in pets, but not extensively researched.

These side effects have been observed in humans on this drug:

- Allergic reaction (labored breathing, hives, etc)
- Vomiting
- Diarrhea
- Skin rash
- Loss of appetite
- Dark urine or jaundice due to liver failure
- Pale tongue, gums, and nose

Fluconazole will react with these drugs:

- Amphotericin B
- Cyclosporine
- Hydrochlorothiazide
- Rifampin

DO NOT GIVE FLUCONAZOLE TO PREGNANT PETS. USE CAUTION WHEN ADMINISTERING THIS DRUG TO DIABETIC PETS, LACTATING PETS, KIDNEY DISEASE, OR LIVER DISEASE. AS THIS DRUG AFFECTS LIVER AND KIDNEYS IN A NEGATIVE WAY, THUS FURTHER DAMAGES THE ORGANS.

# PAIN RELIEF (anti inflammatory drug)

## \* Metecam or Meloxicam ELOXIORAL® Drug

Generic name **Metecam** or **Meloxicam**

**Oral Suspension for dogs and cats - as a anti inflammatory**

Each mL contains 0.5 mg meloxicam in a yellowish suspension with an odour of honey and 1.5 mg of sodium benzoate (Ph. Eur.) as the preservative. Is a nonsteroidal anti-inflammatory drug (NSAID). Meloxicam works by reducing hormones that cause inflammation and pain in the body. It is often given after surgery. **Warnings:** Should not be administered if gastric or intestinal ulceration or bleeding is suspected; if there is evidence of cardiac, hepatic or renal disease; or if there is evidence of a haemorrhagic disorder or individual hypersensitivity to the product. Do not administer concurrently, other steroidal or nonsteroidal anti-inflammatory drugs (NSAIDs), aminoglycoside antibiotics or anticoagulant agents. Pre-treatment with other steroidal or nonsteroidal anti-inflammatory drugs (NSAIDs) may result in additional or increased adverse reactions and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before commencement of treatment depending on the pharmacokinetic properties of the products used previously. Repeat dosing in cats at doses higher than currently recommended has been associated with acute renal failure and death.

## SIDE EFFECTS

Typical adverse reactions of NSAIDs, such as loss of appetite, vomiting, diarrhea, apathy and renal failure have occasionally been reported. These adverse effects are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal. In some cases, death has been associated with some of these adverse reactions.

**Systemic:** anorexia, lethargy, death.

**Gastrointestinal:** vomiting, diarrhea.

**Renal and urinary:** renal disorder, renal failure, urine abnormalities. Repeat dosing in cats at doses higher than currently recommended has been associated with acute renal failure and death. ~ **The most common adverse reactions reported involve the kidneys and the gastrointestinal tract.** Typical symptoms include loss of appetite, depression and vomiting.

It is important in these situations to discontinue treatment immediately and contact your veterinarian. In most cases, the adverse reactions are transient and disappear after termination of treatment but in rare instances may be serious especially if treatment is not discontinued. Consult your veterinarian.

**More frequently reported side effects include:** abdominal pain, anemia, and edema. See below for a comprehensive list of adverse effects. Along with its needed effects, meloxicam may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention.

## Others are

- Arm, back, or jaw pain
- bleeding gums
- **bloating**
- blood in the urine
- blurred vision
- burning upper abdominal or stomach pain
- **canker sores**
- chest tightness or heaviness
- chills
- cloudy urine
- cough
- cramping
- dark urine
- decreased frequency or amount of urine
- difficult or labored breathing
- dilated neck veins
- **dizziness**
- dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position
- general tiredness and weakness
- **headache**
- **hives** or **welts**
- increased blood pressure
- increased sensitivity of the skin to sunlight
- increased thirst
- irregular breathing
- itching, redness, or other discoloration of the skin
- large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs
- light-colored stools
- loss of appetite
- lower side or **back pain**
- noisy breathing
- pain or discomfort in the arms, jaw, back, or neck
- painful or difficult urination
- pains in the stomach, side, or abdomen, possibly radiating to the back
- pinpoint red or purple spots on the skin
- pounding in the ears
- redness, soreness, or itching skin

- **seizures**
- severe and continuing **nausea**
- severe sunburn
- shakiness in the legs, arms, hands, or feet
- skin blisters
- **sore throat**
- sores, ulcers, or white spots on the lips or tongue or inside the mouth
- sores, welting, or blisters
- stomach bloating, burning, cramping, tenderness, or pain
- sweating
- swelling or puffiness of the face
- swollen glands
- trembling or shaking of the hands or feet
- trouble breathing
- unusual bleeding or bruising
- upper right abdominal or stomach pain
- watery or bloody **diarrhea**
- weight gain or loss
- yellow eyes or skin
- Area rash
- blistering, peeling, or loosening of the skin
- bloody or black, tarry stools
- clay-colored stools
- cold, clammy skin
- continuing **vomiting**
- cough or hoarseness
- cracks in the skin
- difficulty with swallowing
- fast, weak pulse
- fever with or without chills
- greatly decreased frequency of urination or amount of urine
- joint or muscle pain
- light headedness
- loss of heat from the body
- puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue
- red skin lesions, often with a purple center
- red, irritated eyes
- red, swollen skin
- scaly skin
- severe stomach pain

- tightness in the chest
  - unpleasant breath odor
  - unusual tiredness or weakness
  - vomiting of blood or material that looks like coffee grounds
- 
- double vision
  - inability to move the arms, legs, or facial muscles
  - inability to speak
  - pains in the chest, groin or legs, especially the calves
  - severe headaches of sudden onset
  - slow speech
  - sudden loss of coordination
  - sudden onset of shortness of breath for no apparent reason
  - sudden onset of slurred speech
  - sudden vision changes

*Get emergency help immediately if any of the following symptoms of overdose occur while taking meloxicam:* **Symptoms of Overdose**

- Blue lips, fingernails, or skin
- convulsions (seizures)
- pain in the chest, upper stomach, or throat
- severe stomach pain
- **skin rash**
- slow or **fast heartbeat**
- swelling around the eyes, face, lips, or tongue
- weight gain (rapid)

Some side effects of meloxicam may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects.

## More Common

**Diarrhea**

**gas**

**heartburn**

**indigestion**

**Abdominal or stomach pain**

**abnormal dreaming**

**anxiety**

**appetite increased**

**bad, unusual, or unpleasant after taste**

- **belching**
- bloated or full feeling
- burning feeling in the chest or stomach
- burning, crawling, itching, **numbness**, prickling, "pins and needles", or tingling feelings
- burning, dry, or itching eyes
- change in taste
- changes in vision
- confusion
- **constipation**
- continuing ringing or buzzing or other unexplained noise in the ears
- decreased urination
- discharge
- dry mouth
- excess air or gas in the stomach
- excessive tearing
- feeling of constant movement of self or surroundings
- feeling sad or empty
- general feeling of discomfort or illness
- **hair loss**
- **hearing loss**
- irritability
- loss of interest or pleasure
- nausea or vomiting
- nervousness
- pain or burning in the throat
- rapid breathing
- redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid
- sensation of spinning
- sleepiness
- **stomach upset**
- sunken eyes
- tenderness in the stomach area
- thirst
- tiredness
- trouble concentrating
- trouble sleeping

# PAIN RELIEF DRUGS

## \* TRAMADOL<sup>®</sup> Drug Used as a Pain relief drug

This medication is used to help relieve moderate to moderately severe pain. Tramadol is similar to opioid (narcotic) analgesics. It works in the brain to change how your body feels and responds to pain.

**SIDE EFFECTS** - Stop using tramadol and call your veterinarian at once if your pet has any of these serious side effects: *seizure; a red, blistering, peeling skin rash; or shallow breathing.* ... Less serious side effects may include: *drowsiness, weakness; vomiting, constipation, loss of appetite; blurred vision; insomnia.*

## Buprenorphine (Buprenex)

Opiate painkiller given to cats to relieve pain from injuries, trauma, and surgeries, as well as for suppressing chronic pain like arthritis <sup>[1]</sup>.

**It is a synthetic partial opiate, around 30 times more potent than morphine**

### Negative symptoms

Fluctuations in body temperature and heart rate

Gastrointestinal problems like loss of appetite, indigestion, constipation, diarrhea, vomiting

Shallow breathing (respiratory depression)

Drop in blood pressure (weakness and lack of activity in your cat)

### It is usually not prescribed to cats with :

Known allergy or hypersensitivity to buprenorphine

Addison's disease, kidney disease, or a compromised thyroid gland

Cardiovascular disease or a head trauma

Liver disease

# PAIN AND INFLAMMATION DRUG

## FIROXIB® Drug

**(use for pain and inflammation associated with osteoarthritis)**

Firocoxib (Previcox) is used in dogs for pain and inflammation associated with osteoarthritis.

Firocoxib is also used to help control postoperative pain and inflammation associated with soft-tissue and orthopedic surgery.

**Side Effects** Firocoxib (Previcox) like other NSAIDs may cause some side effects. The most common side effects of Firocoxib *involve digestive issues such as vomiting and decrease appetite.*

### Other side effects of Firocoxib include:

- Change in bowel movements (diarrhea, or black, tarry or bloody stools)
- Change in behavior (decreased or increased activity level, incoordination, seizure, or aggression)
- Jaundice (yellowing of gums, skin or whites of eyes)
- Increase water consumption or urination changes (frequency, color, or smell)
- Skin irritation (redness, scabs, or scratching)
- Unexpected weight loss

It is important to stop the medication and immediately contact your veterinarian if you think your dog has any medical problems or side effects while taking Firocoxib (Previcox).

\* **Precautions** - Firocoxib (Previcox) is for use in dogs only. Firocoxib should not be given to dogs that are hypersensitive to NSAIDs. Firocoxib should not be given with any other NSAIDs, including: Carprofen (Rimadyl), Meloxicam (Metacam), Etodolac (Etogesic), Deracoxib (Deramaxx), Aspirin. Dogs that are dehydrated, on concomitant diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction are at a greater risk for experiencing adverse events. Use of Firocoxib (Previcox) at doses above the recommended dose in puppies less than seven months of age has been associated with serious adverse reaction, including death. **Drug Interactions** Firocoxib (Previcox) should not be given with other NSAIDs or corticosteroids (e.g., prednisone, cortisone, dexamethasone or tramcinolone).

**Signs of Toxicity** Firocoxib (Previcox) can cause: Loss of appetite, Vomiting, Diarrhea, Dark or tarry stool, Increased thirst, Increase urination, Pale gums, Jaundice, Lethargy, Fast or heavy breathing, Incoordination, Seizures, Behavior changes.

\* *If you suspect or know your dog has had an overdose, it could be fatal so please contact your veterinarian, an emergency vet clinic, or the Pet Poison Helpline at (855) 213-6680 immediately.*

# DRUGS FOR STOMACH ULCERS – ANTACID

## SOMEPRAZOLE® Drug

Omeprazole is an antacid drug commonly used in human medicine.

Excess of stomach acid can lead to gastric ulcers in dogs. As the excess gastric acid can cause reflux, this also poses a threat to the throat areas. There is some research to suggest that administering omeprazole can lead to 'quantitative changes in [gastrointestinal] microbiota of healthy dogs or cats (or human ant acid drugs) "[3]

**Therefore, they no longer have a healthy gut flora defense system to protect from getting infections.**

**Side effects requiring immediate medical attention.**

Along with its needed effects, omeprazole may cause some unwanted effects. Contact your doctor or vet immediately if any of the following side effects occur while taking omeprazole:

- Back, leg, or stomach pain
- bleeding or crusting sores on the lips
- blisters
- bloody or cloudy urine
- chills
- continuing ulcers or sores in the mouth
- difficult, burning, or painful urination
- fever
- frequent urge to urinate
- general feeling of discomfort or illness
- itching, skin rash
- joint pain
- loss of appetite
- muscle aches or cramps
- pain
- red or irritated eyes
- redness, tenderness, itching, burning, or peeling of the skin
- sore throat
- sores, ulcers, or white spots on the lips, in the mouth, or on the genitals
- unusual bleeding or bruising
- unusual tiredness or weakness
  
- Blistering, peeling, or loosening of the skin
- drowsiness

- fast, racing, or uneven heartbeat
- mood or mental changes
- muscle spasms (tetany) or twitching seizures
- nausea
- red skin lesions, often with a purple center
- red, irritated eyes
- trembling
- vomiting

## SYNTHETIC STEROIDS

### \* PREDNISONE® Drug (Prednisolone)

Brand names: Prednisone Intensol and Rayos      **Cortisone (hydrocortisone)**

**Brand names:** A-Hydrocort, Solu-CORTEE, and Cortef.

**What this drug does** -Prednisone and hydrocortisone are synthetic corticosteroids used for suppressing the immune system and inflammation. Both drugs are used to treat many conditions including arthritis, colitis, asthma, bronchitis, skin problems, and allergies.

- Brand names of prednisone include - Prednisone Intensol and Rayos.  
Brand names of hydrocortisone include A-Hydrocort, Solu-CORTEE, and Cortef.
- Common side effects of both prednisone and hydrocortisone include retention of sodium (salt) and fluid, weight gain, high blood pressure (hypertension), potassium loss, headache, muscle weakness, nausea, vomiting, acne, thinning skin, restlessness, and psychiatric disturbances (depression, euphoria, insomnia, mood swings, personality changes, and psychotic behavior).
- **Prolonged use** of prednisone or hydrocortisone can depress the ability of body's adrenal glands to produce corticosteroids.
- **Abruptly stopping prednisone or hydrocortisone** can cause symptoms of corticosteroid insufficiency, **such as nausea, vomiting, weakness, fatigue, decreased appetite, weight loss, diarrhea, abdominal pain, and shock.**
- Prednisone and prednisolone are medications acts on almost every part of the body and have a wide range of effects including reducing inflammation, suppressing the immune system, inhibiting healing, altering mood, stimulating appetite, increasing the secretion of gastric acid, weakening muscles, thinning the skin, and more. In your pet's liver, prednisone is converted to prednisolone.

**Pets with liver problems** are not able to make this conversion effectively, and many veterinarians believe that these pets should only be given prednisolone. Cats also have a limited ability to convert prednisone into prednisolone, so prednisolone is the preferred medication in this species.

## Side effects

### Prednisolone drug (cortisone)

Side Effects of Prednisone and Prednisolone.

When pets have to be on prednisone or prednisolone at high doses and/or for long periods of time the risk of significant side effects like the following rises.

- An increased risk of infection
- Altered behavior, including aggression
- Inhibited growth in young pets
- Development or worsening of diabetes
- Cushing's disease
- Vomiting
- Diarrhea
- Ulceration of the digestive tract
- Lethargy
- Delayed healing

A pet that has been on years of prednisolone will need to have his/her immune system treated and repaired from an initially low immune system (allergy reactions) and further damage from overuse of prednisolone drug injections.

# HEART DRUGS

## Vetmedin prescribed

### PIMOBENDAN® Drug

is the generic name for a heart medication indicated to help dogs suffering from heart failure. This medication makes it easier for the heart to pump blood throughout a dog's body by dilating the blood vessels and decreasing blood pressure.

Pimobendan is administered as a chewable tablet given approximately one hour before feeding, or as directed by a veterinarian. Dog owners should be aware of the potential side effects of pimobendan before giving this medication to their dogs.

#### Side Effects of Pimobendan chemical drug

**Upset Stomach** - the most common side effects of pimobendan are *upset stomach symptoms. These symptoms can include vomiting, diarrhea or loss of appetite.* In fact, approximately one-third of dogs treated with this medication during clinical trials developed poor appetite or diarrhea side effects, according to Drugs.com. Affected dogs may vomit after eating,

**Lethargy or Fainting** - Treatment with this medication can induce lethargy or fatigue side effects in certain dogs, the

**Weakness or Poor Coordination** - A dog treated with pimobendan can appear abnormally weak or uncoordinated. Affected dogs may limit their movements throughout the day, preferring to nap or remain lying down due to sensations of body weakness. A dog can appear unbalanced while walking or running due to poor coordination. These side effects of pimobendan may progressively diminish with continued use of this medication..

**Breathing Difficulties or Cough** - Cough or breathing difficulties can affect dogs taking this medication. In clinical trials for pimobendan, approximately 7 percent of dogs exhibited cough symptoms, Drugs.com reports. A pet owner may notice that his dog begins to wheeze or becomes out of breath quickly due to breathing difficulties.

**Sudden Death**- Approximately 6 percent of dogs treated with pimobendan during clinical trials experienced sudden death, Drugs.com reports. This medication alters the way in which the heart pumps blood through the body, which increases the risk of potentially fatal side effects. Pet owners should be aware of these risks before giving pimobendan to their dogs. If your dog is on heart drugs, search online to find side effects, so you are aware and what to watch out for.

As your Vet wont be warning you or telling your dog is having allergies to the drug or side effects to them.\* See alternatives (or in conjunction) for high blood pressure, heart conditions: e.g

*Magnesium powder or liquid ReMag Magnesium,*

*Homeopathic HBP 148 100ml formula, Homeopathic Congestive Heart 22 drops and herbal heart tonic etc* **See HELP SHEET beside products information.**

# HIGH BLOOD PRESSURE DRUGS

## ENALPRIL MALEATE<sup>®</sup> Drug

"Enalapril is an angiotensin-converting enzyme (ACE) inhibitor that is commonly prescribed by veterinarians for the treatment of high blood pressure in cats and dogs. It can also be used in conjunction with other medications for the treatment of mild to severe heart failure.

### Side Effects and Drug Reactions

- ~ Allergic reaction (labored breathing, hives, etc)
- ~ Vomiting
- ~ Diarrhea
- ~ Lethargy
- ~ High blood pressure
- ~ Fever
- ~ Dizziness
- ~ Fainting
- ~ Ulceration of the digestive tract

**REFERENCE:** READ ABOUT MORE side SERIOUS effects

<https://www.webmd.com/drugs/2/drug-6301-3281/enalapril-maleate-oral/enalapril-oral/details/list-sideeffects>

### ALTERNATIVES

\* See and use possible alternatives options (*or use in conjunction*) for high blood pressure, so your pet is actually being helped to restore balance – rather than control the condition with drugs (*this is why its prescribed ongoing*)

e.g Magnesium liquid or powder, and homeopathic HAMPL HBP 148 100ml formula, Congestive Heart 22 drops and herbal heart tonic, Mitral Valve Insufficiencies See HELP SHEET beside products information.

# URNINARY INCONTINENCE drugs

## PROIN® Drug Chewable tablets (Phenylpropanolamine)

Proin is a veterinarian-prescribed medication for dogs that contains the active ingredient phenylpropanolamine. This medication is indicated to alleviate symptoms associated with urinary incontinence, a condition in which a dog is unable to voluntarily control urine production. Proin works by increasing the tone of the urethral wall and bladder, which helps prevent urine leakage. Dog owners should discuss the side effects of Proin with a veterinarian before giving this medication to their dog.

Restlessness or Irritability - While receiving treatment with Proin, a dog may appear unusually restless or irritable, Side effects of restlessness or irritability may make it difficult for a dog to nap or fall asleep.

Increased Heart Rate or Blood Pressure - Dogs taking Proin may develop an unusually increased heart rate as a side effect of treatment, PetCareRx warns. Certain dogs may also develop hypertension, a condition characterized by increased blood pressure. Diminished

Appetite - Appetite loss may occur in dogs taking Proin,

Allergic Reaction -. Symptoms of a hypersensitive reaction in dogs include weakness, seizures, pale gums, facial swelling and difficulty breathing. Dogs who exhibit any of these side effects after taking a dose of Proin should be transported to an emergency veterinarian clinic immediately. Do not give with ephedrine or epinephrine, or within two weeks of using a MAOi, such as Anipryl, selegiline or a Preventic Collar.

Proin may cause restlessness and hyperactivity, pale gums, difficulty urinating, irritability, hypertension, and decreased appetite.

Hypertension risk increases if Proin is given concurrently with non-steroidal anti-inflammatory drugs (NSAIDs), or with reserpine, tricyclic antidepressants, amitraz (the active ingredient of the Preventic tick control collar and canine Promeris, a flea control product), or ganglionic blocking agents. Conditions such as glaucoma, prostatic hypertrophy, hypertension, hyperthyroidism, diabetes mellitus, and cardiovascular disorders, may worsen

## PROPALIN® Drug - syrup

(phenylpropanolamine in the form of phenylpropanolamine hydrochloride)

Propalin drug is used in cases where urethral sphincter mechanism incompetence - which is the underlying cause of the incontinence.

But it won't get rid of any underlying infection or heal surgery damages after spaying.

### Potential drug Side Effects of Propalin

- Propalin can cause inappetance, loose stools or diarrhoea, restlessness and in severe cases collapse
- Propalin increase blood pressure and heart rate and can cause arrhythmias
- Do not use Propalin in pregnant or lactating animals
- If you suspect side effects then call your vet for advice
- An overdose of Propalin can cause severe cases of the above side effects
- If you suspect an overdose then call your vet as soon as possible
- Wash hands if Propalin comes into contact with the skin

Propalin should be avoided if **Diabetes, Cushings, Liver and Kidney disease as its so toxic to organs.**

Reference: <https://www.petprescription.co.uk/disease-and-illnesses-of-dogs-urinary-incontinence-propalin>

# URINARY TRACT INFECTION

## Urex Drug

Urex (methenamine hippurate) is a urinary tract antiseptic drug used to treat and prevent urinary tract infections. Urex is available in generic form. **Most Common side effects of Urex include:**

- nausea,
- upset stomach,
- vomiting,
- diarrhea,
- abdominal cramps,
- loss of appetite, and skin rash.
- mouth sores,
- headache,
- ringing in the ears,
- muscle cramps, or
- swelling of the arms or leg

## Piroxican (bladder cancer drug)

**Risks and Side Effects of Piroxicam.** Piroxicam can cause some side effects and may react with other medicines in your pet's system. Tell your vet about all other conditions that your pet is suffering from before you begin any treatment program involving piroxicam. Additionally, it's important to notify your vet of any other medicines that your dog may be taking.

**The most common side effects of piroxicam include the following:**

- Stomach and intestinal ulcers and bleeding, dark or tarry stools (from bleeding into the intestines), vomiting, loss of appetite and peritonitis (from intestinal perforation).
- Some of these are life-threatening problems and they can occur relatively suddenly.
- Loss of blood clotting ability
- Liver damage (in extreme doses only)
- Intestinal ulcers
- Dark colored stool
- Intestinal perforation
- Vomiting
- Diarrhea
- Loss of energy

If you notice any of these side effects, it's crucial that you inform your vet immediately.

**Reference:** <https://www.vetinfo.com/piroxicam-for-dogs.html>

# CONVULSION DRUG

## \* GABAPENTIN® Drug

Gabapentin is often used as a treatment for chronic pain in dogs and anti convulsion drug. Gabapentin is a type of analgesia, sold under the brand name Neurontin among others, is an anticonvulsant medication for dogs is the main reason to prescribe this drugs. Or chronic pain. There are some risks in long-term use of gabapentin in dogs, such as an increased risk of cancer.

### More common in young pets

- Aggressive behavior or other behavior problems
- anxiety
- concentration problems and change in school performance
- crying
- depression
- false sense of well-being
- hyperactivity or increase in body movements
- rapidly changing moods
- reacting too quickly, too emotional, or overreacting
- restlessness
- suspiciousness or distrust

### Common

- Clumsiness or unsteadiness
- continuous, uncontrolled, back-and-forth, or rolling eye movements
- Black, tarry stools
- chest pain
- chills
- cough
- depression, irritability, or other mood or mental changes
- fever
- loss of memory
- pain or swelling in the arms or legs
- **painful or difficult urination**
- shortness of breath
- sore throat
- sores, ulcers, or white spots on the lips or in the mouth
- swollen glands
- unusual bleeding or bruising

- unusual tiredness or weakness
- Abdominal or stomach pain
- blistering, peeling, or loosening of the skin
- clay-colored stools
- **coma**
- confusion, headache
- **convulsions**
- dark urine
- decreased urine output
- diarrhea
- dizziness
- **fast or irregular heartbeat**
- increased thirst
- itching or skin rash
- joint pain
- large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet,
- **loss of appetite**
- muscle ache or pain
- nausea
- red skin lesions, often with a purple center
- red, irritated eyes
- unpleasant breath odor
- **vomiting of blood**
- yellow eyes or skin

**Some side effects of gabapentin may occur that usually do not need medical attention.** These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

**NOTE:** people that have taken their pet to emergency hospital from side effects of drugs, not one vet has acknowledged that acute symptoms or chronic ill health was from the drug given...

- Blurred vision
- cold or flu-like symptoms
- delusions
- dementia
- hoarseness
- lack or loss of strength
- lower back or side pain
- swelling of the hands, feet, or lower legs
- trembling or shaking

- Accidental injury
  - appetite increased
  - back pain
  - bloated or full feeling
  - body aches or pain
  - burning, dry, or itching eyes
  - change in vision
  - change in walking and balance
  - clumsiness or unsteadiness
  - congestion
  - constipation
  - cough producing mucus
- 
- difficulty with breathing
  - dryness of the mouth or throat
  - earache
  - excess air or gas in the stomach or intestines
  - excessive tearing
  - eye discharge
  - feeling faint, dizzy, or lightheadedness
  - feeling of warmth or heat
  - flushed, dry skin
  - flushing or redness of the skin, especially on the face and neck
  - frequent urination
  - fruit-like breath odor
  - impaired vision
  - incoordination
  - increased hunger
  - increased sensitivity to pain
  - increased sensitivity to touch
  - increased thirst
  - indigestion
  - noise in the ears
  - pain, redness, rash, swelling, or bleeding where the skin is rubbed off
  - passing gas
  - redness or swelling in the ear
  - redness, pain, swelling of the eye, eyelid, or inner lining of the eyelid
  - runny nose
  - sneezing
  - sweating

- tender, swollen glands in the neck
- tightness in the chest
- tingling in the hands and feet
- trouble sleeping
- trouble swallowing
- trouble thinking
- twitching
- unexplained weight loss
- voice changes
- vomiting
- weakness or loss of strength
- weight gain

Reference: <https://www.drugs.com/sfx/gabapentin-side-effects.html>

## We just don't know how these poisons get approved by the TGA, FDA and APVMA

There are many many more drugs that are available **and many drugs are re-branded with a new name... people think it's a new drug when its not.**

If you dont see your drug (liquid, oral tablet, or injections, topical applications, skin treatments etc).If you need help, please save a tiny sample and post to clinic with name of drug and your full name and address so we can help -post back to a homeopathic drug detox drops from short term or long term side effects and organs damages.

When we give this drug in homeopathic form in a 30Cor 200C (called tautopathic therapy).

# Snail Bate Poisoning

FOR ALL POISONINGS - Start giving your pet VITAMIN C [Page 77](#)

It is also very important to start Vitamin C therapy as soon as possible. See home therapy page about dosing with Vitamin C. Plus **Activated charcoal**, to absorb poison in the system. [Page 75-76](#)

Prevention is the key

It is safest to avoid the use of all slug and snail bait if you have a pet. If your council allows it, consider chickens and ducks as they are great natural snail and slug controllers! If you do need to use slug and snail bait ALWAYS keep it out of your pet's **reach (remember that dogs, in particular, are good at seeking it out)**. And finally, be careful if you are visiting a friend's or neighbour's garden with your pet as you might not realise they have used snail bait. The warm season brings about plenty of hazards for our pets and one, in particular, is the use of snail and slug bait in the garden. There is a lot of misinformation out there including the idea that there are brands available that are safe for pets. This is not the case and these 'pet-friendly' products can still be dangerous.

**Is snail bait poisonous to dogs?**

Absolutely!

Ingestion can be fatal and it's important to know what to watch out for and how you can prevent poisoning. Dogs seem to be more prone as they are scavengers and love to eat things they shouldn't. Cats can be a bit pickier when it comes to what they eat but are still at risk.

**There are 3 types of snail baits:**

- Metaldehyde- green pellets
- Methiocarb - blue pellets
- Iron EDTA (Multiguard)- brown/yellow pellets

example:

1. Metaldehyde – green pellets
2. Methiocarb – blue pellets

These are the most dangerous and they act on the nervous system causing increased stimulation and can be rapidly fatal if immediate veterinary treatment is not given. While the Multiguard (below) contain a bitterant but these act **ONLY as a deterrent** and many dogs will still eat these pellets.

1. Iron EDTA (Multiguard) – brown/yellow pellets

These are also marketed as 'pet-friendly' & are particularly dangerous if ingested in large amounts.

The iron ingestion may cause gastrointestinal issues such as vomiting and diarrhoea but if enough is eaten, the iron can also cause damage to the liver, spleen, heart, kidneys or brain and even lead to organ failure and death. If your pet ingests this type of slug or snail bait, it's best to seek veterinary advice.

### **The main signs of snail bait poisoning include:**

The notes below relate to metaldehyde and methiocarb toxicity.

### **Signs**

- Excessive drooling
- Depression or restlessness
- Rapid heart rate & panting
- Vomiting & diarrhoea
- Muscle tremors
- Seizures
- Death if not treated
- Animals that survive may develop **liver failure days later**

### **Diagnosis**

- Based on history of ingestion or owners finding eaten packaging.
- Clinical signs suggestive of poisoning
- Blood tests may be performed to assess organ damage and to rule out other causes

### ***What to do if your dog eats snail bait***

If your pet has ingested snail bait (or even if you just think there is a slight possibility your pet has ingested snail bait), you should seek veterinary advice immediately.

Bring the packaging with you if you can so your veterinarian knows what was ingested.

The sooner your pet visits a veterinarian, the better the outcome is likely to be. Your pet will need to be made to vomit to help remove as much of the contents from the stomach as possible.

In some cases, your pet may require and an anaesthetic so that the gastric contents can be washed out via a stomach flush and an enema given to reduce toxicity as much as possible. Supportive care in the form of intravenous fluids may be needed as well as medications to help with muscle tremors and seizures.

**Snail baits can come in the form of pellets, granules, liquids or powders.**

*The poison is mixed with molasses, apple, rice, soybeans or oats. This mixture is what makes it palatable to snails, but it is also what makes it tasty for dogs and cats.*

Even so called "pet friendly" products are dangerous for animals.

Metaldehyde and methiocarb act on the animals **nervous system** causing increased stimulation, and can be fatal if immediate veterinary treatment is not given.

**Multiguard** is less toxic than the other 2 snail baits, but iron toxicity can still be fatal and therefore treatment is recommended even if a small amount has been eaten. Iron toxicity can cause ***gastrointestinal signs*** such as vomiting and diarrhoea, or may cause ***damage to the liver, spleen, heart, kidneys or brain.***

### **Conventional Treatment**

Supportive care including fluid therapy is given. In most cases the animal will require a gastric lavage (stomach flush) and an enema under general anaesthesia.

### **Prognosis**

If prompt, aggressive treatment is given, the prognosis is good. There can be varying factors that may also influence recovery, including current health of animal, age, amount of poison ingested and time when treatment is given.

Signs may last for a number of days, and hospitalisation will be required for severe poisoning.

### **Follow up and aftercare**

Animals may go home with a course of **activated charcoal**, which helps to absorb any remaining poison in the system. Repeat blood tests may be performed to check for any liver damage.

### ***Remember***

**Don't use snail baits if you have pets**

## When "detoxing" use in conjunction with

### "Activated Charcoal"

#### HEAVY METAL POISONINGS (e.g. vaccines)

#### INGESTED POISONS ... absorb poisons in the system

Use activated charcoal powder first you may see a big result quickly.

\* **SUGGESTION** (*double the amount if poisoned with heavy metals*)

Buy on-line to find on line stores that sell it or find at health food stores.

Health Benefits The American Journal of Gastroenterology published a study reporting a scientific claim that activated charcoal can reduce the build-up of **nitrogen waste** in the body; e.g. ..UREA levels (toxic waste) from kidneys.

Research studies have also shown that **activated charcoal decreases the toxic load on the body by over 60% with daily use.** This is due to its ability to bind and grab up chlorine, pesticides and herbicides, bacteria, viruses, yeast, parasites, toxic chemicals and endotoxic metabolites.

Endotoxins These are substances that are produced within our body. This would include toxins produced by our gut bacteria, molecules formed by incomplete liver conjugation and ammonia which can arise due to inadequate kidney filtration.

Kidney Cleansing-The kidneys are critical to filtering and removing toxins through the urinary system. Additionally, it takes urea out of the body and when it is functioning poorly it can lead to elevated ammonia, which is extremely toxic to the brain. Poorly functioning kidneys are also major contributors to blood pressure problems and cardiovascular disease. It also absorbs toxins in the kidneys and livers preventing a toxic overload, and at the same time allowing the vital organs (.. liver and kidneys) in the body to function normally. It also can help to eliminate the toxins in the body because it stimulates the bile flow

## SUGGESTION

to assist your pet with – whether acute or chronic kidney weakness. Starting adding activated charcoal to your pets daily meals. e.g. get the "activated charcoal" in powder form.

**PUPPY** give approx: 1/4 teaspoon of the powder to meals or milk drinks.

**ADULT PETS:** use 1 capsule of the powder or 1/2 a teaspoon in meals and/or milk drinks.

**MULTIPLE PETS** (cats and dogs), add 2 capsules or 1 teaspoon the powder into a jar with a screw lid (one or 2 cups of filtered water). Can add the homeopathic detox remedies (along with) things like the Bioplasma drops (tissue salts) is great for growing infants and run down pets and kittens who may have had parents who were mal-nutrition or feral etc, homeopathic **ParaExpel wormer** and **DrawingOut 16 drops** if teeth or skin issues. Shake before adding to each meal or/milk drinks. pour about a tablespoon or so of the liquid each time.

**TWICE a day for one month** to help detox from the heavy metals from the vaccines.

**Once pet has recovered, continuing using the Activiated Charcoal powder once a day for good health.**

For humans a capsule in the evening before bed. Dr Robin Murphy recommends.

## ***Fast Anti- doting poisonous drug/ vaccine side effects***

### *Oral Vitamin C*

**Use for Emergency FOR FAST – “ANTIDOTE of the poisons” e.g. Vaccination side effects, and other toxic pet products. We use a water-soluble type of Vitamin C - that gets into the blood stream straight away.**

Using SODIUM ASCORBATE Vitamin C POWDER *\* you can be bought from health food stores and on the internet. also buy some small oral syringes.*

#### **A Guideline Oral Dosing**

**HORSE / COW / SHEEP** - ORAL dosing Sodium Ascorbate vitamin C crystals, dissolve 5 teaspoons in 1 cup of filtered water, oral syringe approx. 20ml, small sips at a time. repeating every 2 to 3 hours

**DOG / KID / SWINE / DOE** - ORAL dosing Sodium Ascorbate vitamin C crystals, dissolve 5 teaspoons in 1 cup of filtered water, then using an oral syringe - syringe small sips at a time approx. 15ml repeating every 2 to 3 hours

**CAT / Toy Dog / or Other small animals** - ORAL Vit C - Sodium Ascorbate vitamin C crystals, dissolve 3 teaspoons in 1 cup of filtered water. then using an oral syringe, syringe side of mouth small sips at a time until approx. 5ml has been taken - repeating every 2 to 3 hours

**Kitten (infant small animals)** - ORAL Vit C - Sodium Ascorbate vitamin C crystals, dissolve 3 teaspoons in 1 cup of filtered water. then using an oral syringe, syringe side of mouth small sips at a time until approx. 2ml has been taken - repeating every 2 to 3 hours

**Continue for at least 4 to 6 oral dosings, a day.** For example.... say a dose every 3 to 4 hours. e.g 8am then or 11am, then 3pm, 6pm and 10pm. A hug improvement will be seen with that day. Repeat each day, then upon animal

improvement can stop or reduce to casual three times a day if you feel still not 100%. The reason why we need to do so many repeats is that vitamin c does not stay in the body for very long.

## Homeopathic "Tautopathic" Treatment

The homeopathic DOXYCYCLINE - broad spectrum antibiotic (Vibramycin®, Doxychel®, Doxy Caps®, Bio-tab®, Monodox®, Doryx®, Doxirobe®) in a 30C or 200C

Will clear side effects as well as in a this will clear symptoms of **Candida yeast infections** or prevent it.

(HAMPL Chemical Detox 110 or HAMPL Pet Cleanse 42 formulas)

If your dog is on one or more chemical drugs, search on-line to find side effects, so you are aware and what to watch out for.

As your Vet won't be warning you or telling your dog is having allergies to the drug or side effects to them.

## The found of Homeopathic medicine the physician Samuel Hahnemann said to "remove the obstacles to the cure

**Dr Robin Murphy ..** says... Allopathic prescribed drugs kill outright 120,000 people per year: these are the reported cases only.

Indirectly they kill many more thousands of people. Modern medicine uses weird numerology to manipulate the statistics and they call it a medical science.

The average American has a 9000% greater chance of dying from allopathic drugs than from a loaded gun in their own home.! A person has double the chance of dying from their doctor than from a car accident.

**Classical Homeopathy is useless in cases of severe pathologies and patients on allopathic drugs.**

My practice consists of 60% to 70% of these types of patients. Most patients today are on 2-3 different drugs. Therefore we should be familiar with the top allopathic drugs being prescribed today. We need to know the top heart, blood pressure and diabetic remedies, etc. We need to know: drug side affects, drug overdoses, withdrawal symptoms, etc. We need to know them as potentized remedies also.

In homeopathic prescribing, I never take a patient off their drugs, therefore the allopaths don't object. I always give low potencies of the indicated remedies along with herbal tinctures. Why? Because low potencies given daily are less prone to be antidoted by the drugs. I give higher potencies for drug overdoses, and acute withdrawal symptoms.

Once the patient is doing better, the drug doses are lowered and eventually stopped. We can get patients off their drugs by treating the underlying reason they are taking the drugs. **As the allopathic drugs are reduced, the cure rate is increased.**

There are 1500 new drugs annually, these drugs are new to the human gene pool. Therefore any approved drug is still one foreign to human DNA for generations. The true toxics effects of any drug may not be known for at least 40 years or longer. Human are being chemically manipulated, poisoned and experimented on, for money and no other reason. Chemotherapy

is as toxic as getting cholera and but we have to pay the chemo poisoning and cholera is free!

We are also being poisoned by Synthetic Chemicals called Xenobiotics that have been around since the early 1900's. Since then there are thousands new xenobiotics every year also. Tap water has 56,000 xenobiotics, the US government tests for only 56 additions of these chemicals added to tap water.

**Since 1900, 2 million pesticides, herbicides and chemicals are in the human and Pets blood stream.** **Example:** In brain tumour cases, a chemical called Sodium Laurel Sulphate may be a cause of the cancer. This chemical is found in most shampoos to create foam, even the ones in the health stores. Sodium Laurel Sulphate is detected in the blood stream after one use only. We shampoo our babies hair with a cancer causing agent on a daily basis!

## The Nine Methods of Tautopathic Prescribing

A practitioner needs to be familiar with the Mims or BNF ( British National Formulary) of allopathic drugs. It is important not to be prejudice about Allopathy, but to understand their drugs, as any morbific agent, such as a grief or trauma. In the treatment of patients who are on drugs, if you can use tautopathy first, you can demonstrate what homeopathy can do.

1. **Tautopathic Prescribing for Drug Side Effects.**

Drug side effects are a major obstacle to the cure. The biggest hazard lies in mixing up the case. We can get acute side effects of drugs and chronic symptoms of the patient confused. It's important to separate these two. We need to learn to distinguish between the side effects of the drugs and the symptoms of the patient when we analyze the case history. We need to learn the symptoms of allopathic drugs like we do with the homeopathic

remedies.

**Example:**

You might ask, "did you have these symptoms before or after a drug?" Then, depending on the answer, you might look at remedies like Oophorinum, Ovarian, Oestrogen etc for hormonal cases.

2. **Tautopathic Prescribing for DRUG LAYERS**

"Never been well since taking a drug or chemical poisonings" is a common phenomenon in modern practice. Drug poisonings can leave a drug layers. If a drug layer exists in the case then it must be removed first. Otherwise the patient's health will not improve. Mixing these drug layers with the general case is a big mistake. Also ignoring the top layer of the case can lead to a failure to cure or even palliate.

3. **Tautopathic Prescribing for DRUG WITHDRAWALS**

The drug withdrawal symptoms are a whole new case. Treat all drug withdrawals as acutes, and use the tautopathic remedies in higher potencies. Flush the drugs out with water, by putting the tautopathic remedies in water.

Also use herbal tinctures to help with the detoxification process. Avena Sativa for morphine or heroin withdrawal. Chemotherapy and radiotherapy should be treated as acute poisonings. Go for the herbs that fit the organ and give the tautopathic remedies.

4. **Tautopathic Prescribing for DRUG Addictions**

Clinical homeopathy can assist greatly in the removal of cravings

and addictions. Both psychological and physiological addictions can be treated effectively. Cravings for cigarettes, alcohol, coffee, marijuana, or allopathic drugs are all treatable with tautopathics. Is this being taught in the homeopathic colleges?

**Example:** for alcohol craving: Lachesis, Quercus rubens, Sulphuric acid or Amethyst, (known as the sobriety stone). However, Alcohol in potency is a great remedy. Alcohol dehydrates the brain and can lead to many chronic disorders, including epilepsy.

5. **Tautopathic Prescribing for DRUG and Chemical OVERDOSES**

Acute drug overdoses are a common cause of death in our society. Knowing the correct tautopathic or homeopathic remedies can save someone's life.

**Example:** Barbiturate overdose symptoms are slurred speech, drowsiness, murmuring, and heavy eyes, followed by coma and death. Phenobarb. in potency is the antidote for phenobarb. The clinical picture of Gelsemium also fits.

6. **Tautopathic Prescribing of ALLOPATHIC Drugs based on their ALLOPATHIC indications, AS IN THE BNF / MIMS.**

In spite of all the new provings going on these days, I believe learning allopathic drugs is more important. We instantly have 1,000 new chemical remedies. This involves prescribing a remedy based on its allopathic indications, because it is already a well-proven remedy.

**Example:** Ritalin is used in hyperactivity therefore it has that similar affinity, for that disease. It is a stimulant that treats over stimulated children. Therefore Ritalin in potency is a great remedy for treating hyperactivity.

**Example:** Dopamine is the drug for Parkinson's disease. Therefore Dopamine is a remedy for Parkinson's disease. These patients also have a dopamine deficiency.

**Example:** a child is a Silica constitution at birth, but after abuse goes into a Staphisagria layer, then, after difficult teenage years he gets given Prozac.

These remedies have an affinity for each other, the fact that prozac treats a Staphisagria like condition means that the patient is sensitive to prozac. Drugs like prozac are new remedies: they stand with staphisagria, Arsenicum & Carcinosis.

We must be unprejudiced and treat them like another remedy.

7. **Tautopathic Prescribing of ALLOPATHIC Drugs and Chemicals based on their HOMEOPATHIC indications.**

Remedies like Lithium, Alcohol, Mercury, Lead and Adrenalin have been in the repertory for over a hundred years; via provings and case histories we have the indications for prescribing them as homeopathic remedies. That means any drug or chemical can be used as a remedy to treat things beyond it's normal allopathic indications.

8. **Tautopathic Prescribing of PREVENTATIVE REMEDIES**

Also known as prophylactic tautopathy: If one knows a person is about to be exposed to a toxic substance then one can give a tautopathic remedy to help them through the toxic exposure.

**Example:** Before a visit to the dentist give the anaesthetic in potency, especially if the patient has a history of poor recovery from that drug. Likewise for chemotherapy you can give Cadmium sulph., on the day of the chemo treatment to lessen the side effects of the treatment.